

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 903 349 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

24.03.1999 Bulletin 1999/12

(21) Application number: 98114971.9

(22) Date of filing: 10.08.1998

(51) Int. Cl.⁶: **C07D 295/12**, C07D 207/09,
C07D 207/20, C07D 209/42,
C07D 211/16, C07D 211/18,
C07D 213/81, C07D 215/48,
C07D 215/54, C07D 307/68,
C07D 307/84, C07D 317/68,
C07D 333/34, C07D 333/38,
C07D 333/68, C07D 401/12,
C07D 409/12, A61K 31/445,
A61K 31/495

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

Designated Extension States:

AL LT LV MK RO SI

(30) Priority: 18.08.1997 US 56001 P

(71) Applicant:

F. HOFFMANN-LA ROCHE AG
4070 Basel (CH)

(72) Inventors:

- Gong, Leyi
San Mateo, California 94404 (US)
- Kertesz, Denis John
Mountain View, California 94043 (US)
- Smith, David Bernard
San Mateo, California 94403 (US)

- Talamas, Francisco Xavier
San Carlos, California 94070 (US)
- Wilhelm, Robert Stephen
Los Altos, California 94022 (US)


(74) Representative: Braun, Axel et al
F.Hoffmann-La Roche AG
Patent Department (PLP),
124 Grenzacherstrasse
4070 Basel (CH)

Remarks:

A request for correction of the description has been filed pursuant to Rule 88 EPC. A decision on the request will be taken during the proceedings before the Examining Division (Guidelines for Examination in the EPO, A-V, 3.).

(54) **CCR-3 receptor antagonists**

(57) This invention relates to certain cyclic amine derivatives of Formula (I)



that are CCR-3 receptor antagonists, pharmaceutical compositions containing them, methods for their use and methods for preparing these compounds.

EP 0 903 349 A2

Description

[0001] This invention relates to certain cyclic amine derivatives that are CCR-3 receptor antagonists, pharmaceutical compositions containing them, methods for their use and methods for preparing these compounds.

[0002] Tissue eosinophilia is a feature of a number of pathological conditions such as asthma, rhinitis, eczema, inflammatory bowel diseases and parasitic infections ((see Bousquet, J. et al. *N. Eng. J. Med.* 323: 1033-1039 (1990) and Kay, A.B. and Corrigan, C.J. *Br. Med. Bull.* 48:51-64 (1992)). In asthma, eosinophil accumulation and activation are associated with damage to bronchial epithelium and hyperresponsiveness to constrictor mediators. It is established that chemokines such as RANTES, eotaxin, MCP-2, MCP-3 and MCP-4 activate eosinophils ((see Baggiolini, M. and Dahinden, C.A. *Immunol. Today* 15:127-133 (1994), Rot, A. M. et al. *J. Exp. Med.* 176, 1489-1495 (1992) and Ponath, P.D. et al. *J. Clin. Invest.*, Vol. 97, #3, 604-612 (1996)). However, unlike RANTES and MCP-3 which also induce the migration of other leukocyte cell types, eotaxin is selectively chemotactic for eosinophils ((see Griffith-Johnson, D.A et al. *Biochem. Biophys. Res. Commun.* 197:1167 (1993) and Jose, P.J. et al. *Biochem. Biophys. Res. Commun.* 207, 788 (1994)). Specific eosinophil accumulation was observed at the site of administration of eotaxin whether by intradermal or intraperitoneal injection or aerosol inhalation ((see Griffith-Johnson, D.A et al. *Biochem. Biophys. Res. Commun.* 197:1167 (1993); Jose, P.J. et al. *J. Exp. Med.* 179, 881-887 (1994); Rothenberg, M.E. et al. *J. Exp. Med.* 181, 1211 (1995) and Ponath, P.D. *J. Clin. Invest.*, Vol. 97, #3, 604-612 (1996)).


[0003] The CCR-3 receptor has been identified as a major chemokine receptor which eosinophils use for their response to eotaxin, RANTES and MCP-3. It is expressed predominantly on the surface of eosinophils and is highly selective for eotaxin. When transfected into a murine pre- β lymphoma line, the CCR-3 receptor bound eotaxin, RANTES and MCP-3 and conferred chemotactic responses on these cells to these chemokines ((see Ponath, P.D. et al. *J. Exp. Med.* 183, 2437-2448 (1996)).

[0004] Recently, studies have shown that pretreatment of eosinophils with an anti-CCR-3 mAb completely inhibits eosinophil chemotaxis to eotaxin, RANTES and MCP-3 ((see Heath H. et al. *J. Clin. Invest.*, Vol. 99, #2, 178-184 (1997)) indicating that CCR-3 antagonists are useful for the treatment of eosinophil-mediated inflammatory diseases.

[0005] Glucocorticoids such as dexamethasone, methprednisolone and hydrocortisone have been used for treating many eosinophil-related disorders, including bronchial asthma ((R. P. Schleimer et. al., *Am. Rev. Respir. Dis.*, 141, 559 (1990)). The glucocorticoids are believed to inhibit IL-5, IL-3 mediated eosinophil survival in these diseases. However, prolonged use of glucocorticoids can lead to side effects such as glaucoma, osteoporosis and growth retardation in the patients ((see Hanania N.A et al., *J. Allergy and Clin. Immunol.*, Vol. 96, 571-579 (1995) and Saha M.T. et al, *Acta Paediatrica*, Vol. 86, #2, 138-142 (1997)). It is therefore desirable to have an alternative means of treating eosinophil related diseases without incurring these undesirable side effects.

[0006] The present invention provides a means of combating eosinophil induced diseases, such as asthma.

[0007] In a first aspect, this invention provides compounds selected from the group of compounds represented by Formula (I):



wherein:

T and U are both nitrogen; or one of T and U is nitrogen and the other is carbon;

R^1 and R^2 are, independently of each other, hydrogen or alkyl;

n is an integer from 0 to 2, provided that when n is 0, either T or U is carbon;

m is an integer from 0 to 3;

Ar and Ar^1 are, independently of each other, aryl or heteroaryl;

F is alkylene, alkenylene or a bond, provided that when T and U are nitrogen and F is alkylene, then R^4 is not aryl.

Each R is independently hydrogen or alkyl, or R together with either R^3 or R^4 and the atoms to which they are

attached form a carbocycle or a heterocycle;

R³ and R⁴ are, independently of each other, selected from:

(i) hydrogen, alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroaralkyl, heterocyclyl, heterocyclalkyl, heteroalkyl, cyano or -(alkylene)-C(O)-Z where Z is alkyl, haloalkyl, alkoxy, haloalkyloxy, hydroxy, amino, mono- or disubstituted amino, aryl, aralkyl, aryloxy, aralkyloxy, heteroaryl, heteroaryloxy or heteroaralkyloxy, provided that both R³ and R⁴ are not hydrogen; or

(ii) R³ and R⁴ together with the carbon atom to which they are attached form a carbocycle or a heterocycle;

E is -C(O)N(R⁵)-, -SO₂N(R⁵)-, -N(R⁶)C(O)N(R⁵)-, -N(R⁶)SO₂N(R⁵)-, -N(R⁶)C(S)N(R⁵)-, -N(R⁶)C(O)-, -N(R⁶)C(O)O-, -OC(O)N(R⁶)- or -N(R⁶)SO₂- wherein:
R⁵ is:

(i) hydrogen, alkyl, acyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, heterocyclalkyl, heteroalkyl, or -(alkylene)-C(O)-Z where Z is alkyl, haloalkyl, alkoxy, haloalkyloxy, hydroxy, amino, mono- or disubstituted amino, aryl, aralkyl, aryloxy, aralkyloxy, heteroaryl, heteroaryloxy or heteroaralkyloxy; or

(ii) R⁵ together with either R³ or R⁴ and the atoms to which they are attached forms a heterocycloamino group; and

R⁶ is hydrogen, alkyl, acyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, heterocyclalkyl, heteroalkyl, or - (alkylene)-C(O)-Z where Z is alkyl, haloalkyl, alkoxy, haloalkyloxy, hydroxy, amino, mono- or disubstituted amino, aryl, aralkyl, aryloxy, aralkyloxy, heteroaryl, heteroaryloxy or heteroaralkyloxy,

provided that when T is nitrogen and E is -C(O)N(R⁵)-, -SO₂N(R⁵)-, -N(R⁶)C(O)N(R⁵)-, -N(R⁶)SO₂N(R⁵)- or -N(R⁶)C(S)N(R⁵)-, then m > 0;

Q is -R⁷-W-R⁸- wherein:

R⁷ is an alkylene chain of between 1-6 carbon atoms inclusive;

R⁸ is a bond or an alkylene chain of between 0-4 carbon atoms inclusive;

W is a bond or a group selected from -C(O)-, -NR⁹-, -O-, -S(O)₀₋₂-, C(O)N(R⁹)-, -N(R⁹)C(O)-, -N(R⁹)SO₂-, -SO₂N(R⁹)-, -N(R⁹)C(O)N(R⁹)-, -N(R⁹)SO₂N(R⁹)- or -N(R⁹)C(S)N(R⁹)- wherein:

R⁹ is hydrogen, alkyl, acyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, heterocycloalkyl, heteroalkyl, or - (alkylene)-C(O)-Z where Z is alkyl, haloalkyl, alkoxy, haloalkyloxy, hydroxy, amino, mono- or disubstituted amino, aryl, aralkyl, aryloxy, aralkyloxy, heteroaryl, heteroaryloxy or heteroaralkyloxy,

provided that when T is nitrogen and U is carbon then W is not -C(O)N(R⁹)-, and prodrugs, individual isomers, mixtures of isomers and pharmaceutically acceptable salts thereof.

[0008] In a second aspect, this invention provides pharmaceutical compositions containing a therapeutically effective amount of a compound of Formula (I) or its pharmaceutically acceptable salt and a pharmaceutically acceptable excipient.

[0009] In a third aspect, this invention provides a method of treatment of a disease in a mammal treatable by administration of a CCR-3 receptor antagonist, comprising administration of a therapeutically effective amount of a compound of Formula (I) or its pharmaceutically acceptable salt. The disease states include respiratory diseases such as asthma.

[0010] Unless otherwise stated, the following terms used in the specification and claims have the meanings given below:

[0011] "Alkyl" means a linear saturated monovalent hydrocarbon radical of one to six carbon atoms or a branched saturated monovalent hydrocarbon radical of three to six carbon atoms, e.g., methyl, ethyl, propyl, 2-propyl, pentyl, and the like.

[0012] "Alkenyl" means a linear monovalent hydrocarbon radical of two to six carbon atoms or a branched monovalent hydrocarbon radical of three to six carbon atoms, containing at least one double bond, e.g., ethenyl, propenyl, and the like.

[0013] "Alkylene" means a linear saturated divalent hydrocarbon radical of one to six carbon atoms or a branched saturated divalent hydrocarbon radical of three to six carbon atoms, e.g., methylene, ethylene, propylene, 2-methylpropylene, pentylene, and the like.

[0014] "Alkenylene" means a linear divalent hydrocarbon radical of two to six carbon atoms or a branched divalent hydrocarbon radical of three to six carbon atoms, containing at least one double bond, e.g., ethenylene, 2,4-pentadienylene, and the like.

[0015] "Acyl" means a radical $-C(O)R$ where R is hydrogen, alkyl, alkenyl, cycloalkyl, heteroalkyl, haloalkyl, aryl, aralkyl, heteroaralkyl or heteroaryl, e.g., acetyl, benzoyl, thenoyl, and the like.

[0016] "Acyloxy" means a radical $-OC(O)R$ where R is hydrogen, alkyl, alkenyl, cycloalkyl, heteroalkyl, haloalkyl or optionally substituted phenyl, e.g., acetoxy, benzoyloxy, and the like.

5 **[0017]** "Acylamino" means a radical $-NRC(O)R'$ where R is hydrogen or alkyl and R' is hydrogen, alkyl, alkenyl, cycloalkyl, heteroalkyl, haloalkyl or optionally substituted phenyl, e.g., acetylamino, trifluoroacetylamino, benzoylamino, methylacetylamino, and the like.

[0018] "Halo" means fluoro, chloro, bromo or iodo, preferably fluoro and chloro.

10 **[0019]** "Haloalkyl" means alkyl substituted with one or more same or different halo atoms, e.g., $-CH_2Cl$, $-CF_3$, $-CH_2CF_3$, $-CH_2CCl_3$, and the like.

[0020] "Cycloalkyl" means a saturated monovalent cyclic hydrocarbon radical of three to six ring carbons, e.g., cyclopropyl, cyclohexyl, and the like.

[0021] "Carbocycle" means a saturated, cyclic group of 3 to 6 ring atoms in which all the ring atoms are carbon, e.g., cyclopentyl, cyclohexyl, and the like.

15 **[0022]** "Monosubstituted-amino" means a radical $-NHR$ where R is alkyl, heteroalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl or optionally substituted phenyl, e.g., methylamino, (1-methylethyl)amino, phenylamino, and the like.

[0023] "Disubstituted-amino" means a radical $-NRR'$ where R and R' are independently alkyl, alkenyl, heteroalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl or optionally substituted phenyl. Representative examples include, but are not limited to, dimethylamino, methylethylamino, di(1-methylethyl)amino, methylbenzylamino, and the like.

20 **[0024]** "Aryl" means a monovalent monocyclic or bicyclic aromatic hydrocarbon radical of 6 to 10 ring atoms, and optionally substituted independently with one or more substituents, preferably one or two substituents selected from alkyl, haloalkyl, heteroalkyl, cycloalkyl, cycloalkylalkyl, halo, cyano, nitro, acyloxy, optionally substituted phenyl, heteroaryl, heteroaralkyl, amino, monosubstituted amino, disubstituted amino, acylamino, hydroxylamino, amidino, guanidino, cyanoguanidinyl, hydrazino, hydrazido, $-OR$ [where R is hydrogen, alkyl, haloalkyl, alkenyl, cycloalkyl, cycloalkylalkyl, optionally substituted phenyl, heteroaryl or heteroaralkyl], $-S(O)_nR$ [where n is an integer from 0 to 2 and R is hydrogen, alkyl, haloalkyl, alkenyl, cycloalkyl, cycloalkylalkyl, optionally substituted phenyl, heteroaryl, heteroaralkyl, amino, mono or disubstituted amino], $-C(O)R$ (where R is hydrogen, alkyl, alkenyl, cycloalkyl, heteroalkyl, haloalkyl or optionally substituted phenyl), $-COOR$ (where R is hydrogen, alkyl, optionally substituted phenyl, heteroaryl or heteroaralkyl), $-(alkylene)COOR$ (where R is hydrogen, alkyl, optionally substituted phenyl, heteroaryl or heteroaralkyl), methylenedioxy, 1,2-ethylenedioxy, $-CONR'R''$ or $-(alkylene)CONR'R''$ (where R' and R'' are independently selected from hydrogen, alkyl, cycloalkyl, haloalkyl, cycloalkylalkyl, optionally substituted phenyl, heteroaryl and heteroaralkyl). More specifically the term aryl includes, but is not limited to, phenyl, 1-naphthyl, 2-naphthyl, and derivatives thereof.

35 **[0025]** "Optionally substituted phenyl" means a phenyl group which is optionally substituted independently with one, two or three substituents selected from alkyl, haloalkyl, halo, nitro, cyano, $-OR$ (where R is hydrogen or alkyl), $-NRR'$ (where R and R' are independently of each other hydrogen or alkyl), $-COOR$ (where R is hydrogen or alkyl) or $-CONR'R''$ (where R' and R'' are independently selected from hydrogen or alkyl).

40 **[0026]** "Heteroaryl" means a monovalent monocyclic or bicyclic aromatic radical of 5 to 10 ring atoms containing one, two, or three ring heteroatoms selected from N, O, or S, the remaining ring atoms being C. The aromatic radical is optionally substituted independently with one or more substituents, preferably one or two substituents selected from alkyl, haloalkyl, heteroalkyl, cycloalkyl, cycloalkylalkyl, halo, cyano, nitro, acyloxy, optionally substituted phenyl, amino, monosubstituted amino, disubstituted amino, acylamino, hydroxyamino, amidino, guanidino, cyanoguanidinyl, hydrazino, hydrazido, $-OR$ [where R is hydrogen, alkyl, haloalkyl, alkenyl, cycloalkyl, cycloalkylalkyl or optionally substituted phenyl], $-S(O)_nR$ [where n is an integer from 0 to 2 and R is hydrogen, alkyl, haloalkyl, alkenyl, cycloalkyl, cycloalkylalkyl, optionally substituted phenyl, amino, mono or disubstituted amino], $-C(O)R$ (where R is hydrogen, alkyl, alkenyl, cycloalkyl, heteroalkyl, haloalkyl or optionally substituted phenyl), $-COOR$ (where R is hydrogen, alkyl, or optionally substituted phenyl), $-(alkylene)COOR$ (where R is hydrogen, alkyl or optionally substituted phenyl), methylenedioxy, 1,2-ethylenedioxy, $-CONR'R''$ or $-(alkylene)CONR'R''$ (where R' and R'' are independently selected from hydrogen, alkyl, cycloalkyl, haloalkyl, cycloalkylalkyl or optionally substituted phenyl). More specifically the term heteroaryl includes, but is not limited to pyridyl, pyrrolyl, thiophene, pyrazolyl, thiazolyl, imidazolyl, pyrimidinyl, thiadiazolyl, indolyl, carbazolyl, azaindolyl, benzofuranyl, benzotriazolyl, benzisoxazolyl, purinyl, quinolinyl, benzopyranyl, and derivatives thereof.

55 **[0027]** "Heterocycloamino" means a saturated or unsaturated monovalent cyclic group of 5 to 8 ring atoms, wherein at least one ring atom is N and optionally contains a second ring heteroatom selected from the group consisting of N, O, or $S(O)_n$ (where n is an integer from 0 to 2), the remaining ring atoms being C. The heterocycloamino ring may be fused to a heteroaryl ring, or it may be optionally substituted independently with one or more substituents, preferably one or two substituents, selected from alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, halo, cyano, acyl, amino, monosubstituted amino, disubstituted amino, $-COOR$ (where R is hydrogen or alkyl), $-XR$

(where X is O or S(O)_n, n is an integer from 0 to 2, and R is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heteroaryl or heteroaralkyl) or -CONR'R" (where R' and R" are independently selected from hydrogen or alkyl). Representative examples include but are not limited to pyrrolidino, piperidino, 4-benzoylpiperidino, morpholino, piperazino, 4-(4-benzyloxyphenyl)piperazino, indolino, and the like.

[0028] "Heterocycle" or "Heterocyclyl" means a saturated or unsaturated cyclic radical of 3 to 8 ring atoms in which one or two ring atoms are heteroatoms selected from N, O, or S(O)_n (where n is an integer from 0 to 2). The heterocycle ring may be optionally substituted independently with one, two or three substituents selected from alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, halo, cyano, acyl, acylamino, amino, monosubstituted amino, disubstituted amino, -COOR (where R is hydrogen or alkyl), -XR (where X is O or S(O)_n, where n is an integer from 0 to 2 and R is hydrogen, alkyl, haloalkyl, cycloalkyl, aralkyl, aryl, heteroaryl or heteroaralkyl) or -CONR'R" (where R' and R" are independently selected from hydrogen or alkyl). Representative examples include, but are not limited to tetrahydropyranyl, piperidino, 1-(4-chlorophenyl)piperidino, and the like.

[0029] "Heteroalkyl" means an alkyl, cycloalkyl, or cycloalkylalkyl radical as defined above, carrying a substituent containing a heteroatom selected from N, O, S(O)_n where n is an integer from 0 to 2. Representative substituents include -NR^aR^b, OR^a or -S(O)_nR^c, wherein n is an integer from 0 to 2, R^a is hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, optionally substituted phenyl, or -COR (where R is alkyl), R^b is hydrogen, alkyl, -SO₂R (where R is alkyl or hydroxyalkyl), -SO₂NRR' (where R and R' are independently of each other hydrogen or alkyl), -CONR'R" (where R' and R" are independently selected from hydrogen or alkyl) and R^c is hydrogen, alkyl, cycloalkyl, cycloalkylalkyl, optionally substituted phenyl, amino, monosubstituted amino, or disubstituted amino. Representative examples include, but are not limited to 2-methoxyethyl, benzyloxymethyl, thiophen-2-ylthiomethyl, and the like.

[0030] "Hydroxyalkyl" means a linear monovalent hydrocarbon radical of two to six carbon atoms or a branched monovalent hydrocarbon radical of three or six carbons substituted with one or two hydroxy groups, provided that if two hydroxy groups are present they are not both on the same carbon atom. Representative examples include, but are not limited to, 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, 1-(hydroxymethyl)-2-methylpropyl, 2-hydroxybutyl, 3-hydroxybutyl, 4-hydroxybutyl, 2,3-dihydroxypropyl, 1-(hydroxymethyl)-2-hydroxyethyl, 2,3-dihydroxybutyl, 3,4-dihydroxybutyl and 2-(hydroxymethyl)-3-hydroxypropyl, preferably 2-hydroxyethyl, 2,3-dihydroxypropyl, and 1-(hydroxymethyl)-2-hydroxyethyl.

[0031] "Cycloalkylalkyl" means a radical -R^aR^b where R^a is an alkylene group and R^b is a cycloalkyl group as defined above e.g., cyclopropylmethyl, cyclohexylpropyl, 3-cyclohexyl-2-methylpropyl, and the like.

[0032] "Aralkyl" means a radical -R^aR^b where R^a is an alkylene group and R^b is an aryl group as defined above e.g., benzyl, phenylethyl, 3-(3-chlorophenyl)-2-methylpentyl, and the like.

[0033] "Heteroaralkyl" means a radical -R^aR^b where R^a is an alkylene group and R^b is a heteroaryl group as defined above e.g., pyridin-3-ylmethyl, 3-(benzofuran-2-yl)propyl, and the like.

[0034] "Heterocyclalkyl" means a radical -R^aR^b where R^a is an alkylene group and R^b is a heterocyclyl group as defined above e.g., tetrahydropyran-2-ylmethyl, 4-methylpiperazin-1-ylethyl, and the like.

[0035] "Alkoxy", "haloalkyloxy", "aryloxy", "heteroaryloxy", "aralkyloxy", or "heteroaralkyloxy" means a radical -OR where R is an alkyl, haloalkyl, aryl, heteroaryl, aralkyl, or heteroaralkyl respectively as defined above e.g., methoxy, phenoxy, pyridin-2-yloxy, benzyloxy, and the like.

[0036] "Optional" or "optionally" means that the subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or circumstance occurs and instances in which it does not. For example, "heterocycle group optionally mono- or di- substituted with an alkyl group" means that the alkyl may but need not be present, and the description includes situations where the heterocycle group is mono- or disubstituted with an alkyl group and situations where the heterocycle group is not substituted with the alkyl group.

[0037] "Amino-protecting group" refers to those organic groups intended to protect nitrogen atoms against undesirable reactions during synthetic procedures e.g., benzyl, benzyloxycarbonyl (CBZ), *t*-butoxycarbonyl (BOC), trifluoroacetyl, and the like.

[0038] Compounds that have the same molecular formula but differ in the nature or sequence of bonding of their atoms or the arrangement of their atoms in space are termed "isomers". Isomers that differ in the arrangement of their atoms in space are termed "stereoisomers". Stereoisomers that are not mirror images of one another are termed "diastereomers" and those that are non-superimposable mirror images of each other are termed "enantiomers". When a compound has an asymmetric center, for example, it is bonded to four different groups, a pair of enantiomers is possible. An enantiomer can be characterized by the absolute configuration of its asymmetric center and is described by the *R*- and *S*-sequencing rules of Cahn and Prelog, or by the manner in which the molecule rotates the plane of polarized light and designated as dextrorotatory or levorotatory (i.e., as (+) or (-)-isomers respectively). A chiral compound can exist as either individual enantiomer or as a mixture thereof. A mixture containing equal proportions of the enantiomers is called a "racemic mixture".

[0039] The compounds of this invention may possess one or more asymmetric centers; such compounds can therefore be produced as individual (*R*)- or (*S*)- stereoisomers or as mixtures thereof. For example, if the R³ and R⁴ substit-

uents in a compound of Formula (I) are different, then the carbon to which they are attached is an asymmetric center and the compound of Formula (I) can exist as an (*R*)- or (*S*)-stereoisomer. Unless indicated otherwise, the description or naming of a particular compound in the specification and claims is intended to include both individual enantiomers and mixtures, racemic or otherwise, thereof. The methods for the determination of stereochemistry and the separation of stereoisomers are well-known in the art (see discussion in Chapter 4 of "Advanced Organic Chemistry", 4th edition

J. March, John Wiley and Sons, New York, 1992).

[0040] A "pharmaceutically acceptable excipient" means an excipient that is useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable, and includes an excipient that is acceptable for veterinary use as well as human pharmaceutical use. "A pharmaceutically acceptable excipient" as used in the specification and claims includes both one and more than one such excipient.

[0041] A "pharmaceutically acceptable salt" of a compound means a salt that is pharmaceutically acceptable and that possesses the desired pharmacological activity of the parent compound. Such salts include:

- (1) acid addition salts, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as acetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethane-disulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid, 4-methylbicyclo[2.2.2]oct-2-ene-1-carboxylic acid, glucoheptonic acid, 4,4'-methylenebis-(3-hydroxy-2-ene-1-carboxylic acid), 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynapthoic acid, salicylic acid, stearic acid, muconic acid, and the like; or
- (2) salts formed when an acidic proton present in the parent compound either is replaced by a metal ion, e.g., an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic base such as ethanolamine, diethanolamine, triethanolamine, tromethamine, *N*-methylglucamine, and the like.

[0042] "Leaving group" has the meaning conventionally associated with it in synthetic organic chemistry i.e., an atom or group capable of being displaced by a nucleophile and includes halogen, alkanesulfonyloxy, arenesulfonyloxy, ester, or amino such as chloro, bromo, iodo, mesyloxy, tosyloxy, trifluorosulfonyloxy, methoxy, *N,O*-dimethylhydroxylamino, and the like.

[0043] "Pro-drugs" means any compound which releases an active parent drug according to Formula (I) *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs of a compound of Formula (I) are prepared by modifying functional groups present in the compound of Formula (I) in such a way that the modifications may be cleaved *in vivo* to release the parent compound. Prodrugs include compounds of Formula (I) wherein a hydroxy, sulfhydryl or amino group in compound (I) is bonded to any group that may be cleaved *in vivo* to regenerate the free hydroxyl, amino, or sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to esters (e.g., acetate, formate, and benzoate derivatives), carbamates (e.g., *N,N*-dimethylaminocarbonyl) of hydroxy functional groups in compounds of Formula (I), and the like.

[0044] "Treating" or "treatment" of a disease includes:

- (1) preventing the disease, i.e. causing the clinical symptoms of the disease not to develop in a mammal that may be exposed to or predisposed to the disease but does not yet experience or display symptoms of the disease,
- (2) inhibiting the disease, i.e., arresting or reducing the development of the disease or its clinical symptoms, or
- (3) relieving the disease, i.e., causing regression of the disease or its clinical symptoms.

[0045] A "therapeutically effective amount" means the amount of a compound that, when administered to a mammal for treating a disease, is sufficient to effect such treatment for the disease. The "therapeutically effective amount" will vary depending on the compound, the disease and its severity and the age, weight, etc., of the mammal to be treated.

[0046] The nomenclature used in this application is generally based on the IUPAC recommendations, e.g.,

a compound of Formula (I) where T and U are nitrogen, n and m are 1, R, R¹, R² and R³ are hydrogen, R⁴ is 1-methylethyl, E is -C(O)NH-, F is a bond, Q is -CH₂-, Ar is phenyl, Ar¹ is 3,4-chlorophenyl and the stereochemistry at the carbon to which R³ and R⁴ are attached is *RS* is named, N-{1(*RS*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}benzamide.

[0047] A compound of Formula (I) where T and U are nitrogen, n and m are 1, R, R¹, R² and R³ are hydrogen, R⁴ is 1,1-dimethylethyl, E is -NHC(O)NH-, F is a bond, Q is -CH₂-, Ar is phenyl, Ar¹ is 3,4-chlorophenyl and the stereochemistry at the carbon to which R³ and R⁴ are attached is *RS* is named, 1-{1(*RS*)-[4-(3,4-dichlorobenzyl)piperazin-1-yl-

methyl]-2,2-dimethylpropyl]-3-phenylurea.


[0048] A compound of Formula (I) where T and U are nitrogen, n and m are 1, R, R¹, R² and R³ are hydrogen, E is -C(O)NR⁵, R⁴ and R⁵ together form a 3-pyrrolino ring, F is a bond, Q is -CH₂-, Ar is 4-methylphenyl, Ar¹ is 3,4-chlorophenyl and the stereochemistry at the carbon to which R³ and R⁴ are attached is *RS* is named, {2-(*RS*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-1-(4-methylbenzoyl)-3-pyrroline.

[0049] A compound of Formula (I) where T is nitrogen, U is carbon, n and m are 1, R, R¹, R² and R³ are hydrogen, R⁴ is 1-methylethyl, E is -C(O)NH-, F is a bond, Q is -CH₂-, Ar is 4-methylphenyl, Ar¹ is 3,4-chlorophenyl and the stereochemistry at the carbon to which R³ and R⁴ are attached is *S* is named, N-{1(*S*)-[4-(3,4-dichlorobenzyl)piperidin-1-yl-methyl]-2-methylpropyl}-4-methylbenzamide.

Representative compounds of this invention are as follows:

I. Representative compounds of Formula (I) where T and U = nitrogen; m and n = 1; R=R¹=R²=R³=hydrogen; Q = -CH₂-; E = -C(O)NH- and other groups are as defined below are:

[0050]



CPD #	Stereo- chem.	Ar	F	R ⁴	Ar ¹	M.Pt. °C	Mass. Spec m/e
1	(R)	4-methylphenyl	bond	1-methylethyl	3,4-dichlorophenyl . 2HCl	232-232.4	
2	(S)	4-methylphenyl	bond	1,1-dimethylethyl	3,4-dichlorophenyl . 2HCl	238-242	
3	(S)	4-methoxyphenyl	bond	1-methylethyl	3,4-dichlorophenyl . 2HCl	222.5-223	
4	(RS)	3,4-methylene- dioxyphenyl	bond	1-methylethyl	3,4-dichlorophenyl		477
5	(RS)	3,4-difluorophenyl	bond	1-methylethyl	3,4-dichlorophenyl		469
6	(RS)	thiophen-2-yl	bond	1-methylethyl	3,4-dichlorophenyl		439

CPD #	Stereo- chem.	Ar	F	R ⁴	Ar ¹	M.Pt. °C	Mass. Spec m/e
7	(RS)	2-pyridyl	bond	1-methylethyl	3,4-dichlorophenyl .2HCl	226.2-229.9	
8	(S)	4-methylphenyl	bond	1-methylethyl	3,4-dichlorophenyl .2HCl	229-229.6	
9	(RS)	4-methylphenyl	bond	1-methylethyl	4-nitrophenyl		424
10	(R)	3-cyanophenyl	bond	1-methylethyl	3,4-dichlorophenyl		458
11	(RS)	4-methylthiophenyl	bond	1-methylethyl	3,4-dichlorophenyl		
12	(RS)	4-acetylphenyl	bond	1-methylethyl	3,4-dichlorophenyl		475
13	(RS)	2-benzofuranyl	bond	1-methylethyl	3,4-dichlorophenyl		473
14	(RS)	4-N,N-dimethyl- aminophenyl	bond	1-methylethyl	3,4-dichlorophenyl		476
15	(RS)	2-indolyl	bond	1-methylethyl	3,4-dichlorophenyl		472
16	(RS)	4-biphenyl	bond	1-methylethyl	3,4-dichlorophenyl		509
17	(RS)	4-methylsulfonyl- phenyl	bond	1-methylethyl	3,4-dichlorophenyl		511
18	(RS)	4-aminophenyl	bond	1-methylethyl	3,4-dichlorophenyl		448
19	(S)	2-naphthyl	bond	1-methylethyl	3,4-dichlorophenyl		483

CPD #	Stereo- chem.	Ar	F	R ⁴	Ar ¹	M.Pt. °C	Mass. Spec m/e
20	(RS)	4-pyridyl	bond	1-methylethyl	3,4-dichlorophenyl . 2HCl	150-153	
21	(RS)	4-ethylenephanyl	bond	1-methylethyl	3,4-dichlorophenyl		459
22	(S)	4-methylphenyl	bond	cyclohexyl	3,4-dichlorophenyl . 2HCl	145.9-148.2	
23	(RS)	4-methylphenyl	bond	<i>n</i> -propyl	3,4-dichlorophenyl . 2HCl	199.5-203.5	
24	(RS)	2-quinoliny	bond	1-methylethyl	3,4-dichlorophenyl		484
25	(RS)	thiophen-2-yl	-CH ₂ -	1-methylethyl	3,4-dichlorophenyl		453
26	(RS)	4-cyanophenyl	bond	1-methylethyl	4-chlorophenyl		424
27	(S)	4-amino-5-chloro-2-methoxyphenyl	bond	1,1-dimethylethyl	3,4-dichlorophenyl . 2HCl	194.8-196.1	
28	(RS)	4-(pyrrol-1-yl)-phenyl	bond	1-methylethyl	3,4-dichlorophenyl		498
29	(RS)	5-nitrofuranyl	bond	1-methylethyl	3,4-dichlorophenyl		468
30	(RS)	5-methoxyindol-2-yl	bond	1-methylethyl	3,4-dichlorophenyl		502
31	(RS)	3,4-methylene-dioxyphenyl	bond	1-methylethyl	4-chlorophenyl		443

CPD #	Stereo-chem.	Ar	F	R ⁴	Ar ¹	M.Pt. °C	Mass. Spec m/e
32	(RS)	phenyl	-(CH ₂) ₂ -	1-methylethyl	3,4-dichlorophenyl		461
33	(S)	4-chlorophenyl	bond	methyl	3,4-dichlorophenyl .2HCl	244-246.5	
34	(RS)	4-chlorophenyl	bond	phenyl	3,4-dichlorophenyl .2HCl	117-183	
35	(R)	4-chlorophenyl	bond	cyclohexyl	3,4-dichlorophenyl	149-155	
36	(RS)	5-chloroindol-2-yl	bond	1-methylethyl	3,4-dichlorophenyl		506
37	(RS)	4-biphenyl	bond	2-methylpropyl	3,4-dichlorophenyl		523
38	(RS)	3-chloro-2-nitro-phenyl	bond	1-methylethyl	3,4-dichlorophenyl		512
39	(RS)	2,4,6-trimethylphenyl	bond	1-methylethyl	3,4-dichlorophenyl		475
40	(RS)	4-chlorophenyl	bond	benzyl	3,4-dichlorophenyl .2HCl	137-142	
41	(RS)	anthraquinon-2-yl	bond	1-methylethyl	3,4-dichlorophenyl		563
42	(RS)	benzothiophen-2-yl	bond	1-methylethyl	4-chlorophenyl		455
43	(RS)	2-phenoxyphenyl	bond	1-methylethyl	3,4-dichlorophenyl		525
44	(RS)	4-methylphenyl	ethenylene	1-methylethyl	3,4-dichlorophenyl		473
45	(RS)	4-methylphenyl	bond	1-methylethyl	4-chlorophenyl		

CPD #	Stereo-chem.	Ar	F	R ⁴	Ar ¹	M.Pt. °C	Mass. Spec m/e
46	(RS)	4-methylphenyl	bond	1-methylethyl	4-nitrophenyl		
47	(RS)	4-methylphenyl	bond	1-methylethyl	3,4-difluorophenyl		
48	(RS)	4-methylphenyl	bond	1,1-dimethylethyl	3,4-dichlorophenyl		
49	(RS)	4-methylphenyl	bond	1-methylethyl	2,3-dichlorophenyl		
50	(RS)	4-methylphenyl	bond	1-methylethyl	3-methyl-4-nitrophenyl		
51	(RS)	4-methylphenyl	bond	1-methylethyl	3-chloro-4-fluorophenyl		
52	(R)	4-methylphenyl	bond	1,1-dimethylethyl	3-chloro-4-fluorophenyl		
53	(R)	4-methylphenyl	bond	1,1-dimethylethyl	3-methylbenzothiophen-2-yl		
54	(R)	4-methylphenyl	bond	1,1-dimethylethyl	1-acetylindol-3-yl		
55	(R)	4-methylphenyl	bond	1,1-dimethylethyl	5-nitrothiophen-3-yl		
56	(R)	quinolin-3-yl	bond	1,1-dimethylethyl	3,4-dichlorophenyl		


and are named as:

2. N-{1(S)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-4-methylbenzamide dihydrochloride salt.

3. N-{1(*S*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methoxybenzamide dihydrochloride salt.
4. N-{1(*RS*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-3,4-methylenedioxybenzamide.
7. N-{1(*RS*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}picolinamide dihydrochloride salt.
- 5 8. N-{1(*S*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide dihydrochloride salt.
19. N-{1(*S*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-2-naphthaleneamide.
25. N-{1(*RS*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-2-(thiophen-2-yl)acetamide.
35. N-{1(*R*)-cyclohexyl-2-[4-(3,4-dichlorobenzyl)piperazin-1-yl]ethyl}4-chloro-benzamide.
37. N-{1(*RS*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-3-methylbutyl}-4-phenylbenzamide.
- 10 44. N-{1(*RS*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-3-(4-methylphenyl)acrylamide.

II. Representative compounds of Formula (I) where T = nitrogen; U = carbon; m and n = 1; $R=R^1=R^2=R^3$ =hydrogen; F = bond; E = -C(O)NH- and other groups are as defined below are:

15 [0051]



CPD #	Stereo -chem	Ar	R ⁴	Q	Ar ¹	M.Pt. °C	Mass. Spec. m/e
57	(S)	4-methylphenyl	1-methylethyl	-CH ₂ -	3,4-dichlorophenyl . HCl	87-90	
58	(S)	4-methylphenyl	1,1-dimethylethyl	-CH ₂ -	3,4-dichlorophenyl . HCl	153.1-154.2	
59	(RS)	4-methylphenyl	1-methylethyl	-CH ₂ NHC(O)-	4-amino-5-chloro-2-methoxyphenyl		515.6

CPD #	Stereo -chem	Ar	R ⁴	Q	Ar ¹	M.Pt. °C	Mass. Spec. m/e
60	(RS)	4-chlorophenyl	1-methylethyl	-CH ₂ -	3,4-dichlorophenyl		467.86
61	(S)	4-chlorophenyl	1,1,-dimethylethyl	-CH ₂ -	3,4-dichlorophenyl	63-68	
62	(RS)	4-methylphenyl	1-methylethyl	-CH ₂ -	4-chlorophenyl . HCl	123-128	
63	(R)	4-(2-acetylamino-ethyl)phenyl	1-methylethyl	-CH ₂ -	3,4-dichlorophenyl . HCl	113.3-113.9	
64	(R)	4-[(2-(R)-amino-3-methylbutyryl-amino)ethyl]phenyl	1-methylethyl	-CH ₂ -	3,4-dichlorophenyl . 2 HCl	191-199	
65	(R)	4-(2-aminoethyl)-phenyl	1,1-dimethylethyl	-CH ₂ -	3,4-dichlorophenyl . 2 HCl	192-198	
66	(R)	4-aminomethyl-phenyl	1,1-dimethylethyl	-CH ₂ -	3,4-dichlorophenyl . 2 HCl		476
67	(R)	quinolin-3-yl	1,1-dimethylethyl	-CH ₂ -	3,4-dichlorophenyl . HCl		534
68	(R)	4-methylphenyl	1,1-dimethylethyl	-CH ₂ -	4-nitrophenyl		
69	(R)	pyridin-2-yl	1,1-dimethylethyl	-CH ₂ -	4-nitrophenyl . HCl		425

CPD #	Stereo -chem	Ar	R ⁴	Q	Ar ¹	M.Pt. °C	Mass. Spec. m/e
70	(R)	pyridin-2-yl	1,1-dimethylethyl	-CH ₂ -	4-nitrophenyl		
71	(R)	5-methylthiophen-2-yl	1-methylethyl	-CH ₂ -	3,4-dichlorophenyl . HCl	45-47.5	
72	(R)	4-(2-aminoethyl)-phenyl	1-methylethyl	-CH ₂ -	3,4-dichlorophenyl . 2 HCl		476
73	(R)	4-methylphenyl	1,1-dimethylethyl	-CH ₂ -	3,4-dichlorophenyl . HCl	153.1-154.2	
74	(R)	4-methylsulfonyl-phenyl	1,1-dimethylethyl	-CH ₂ -	3,4-dichlorophenyl	205-214	
75	(R)	5-methylthiophen-2-yl	1,1-dimethylethyl	-CH ₂ -	3,4-dichlorophenyl . HCl	161-162.5	
76	(R)	4-hydroxymethyl-phenyl	1-methylethyl	-CH ₂ -	3,4-dichlorophenyl		463


and are named as:


57. N-{1(S)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide hydrochloride salt.

59. N-{1(*RS*)-[4-(4-Amino-5-chloro-2-methoxyphenylcarbonylaminomethyl)piperidin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide.

III. Representative compounds of Formula (I) where T and U = nitrogen; m and n = 1; $R=R^1=R^2=R^3$ = hydrogen; Q = -CH₂- and F = bond; E = -C(O)NR⁵; R⁵ together with R⁴ and the atoms to which they are attached = heterocycloamino group and other groups are as defined below are:

[0052]




CPD #	Stereo- chem	Ar		Ar ¹	M.Pt. °C
77	(<i>RS</i>)	4-methylphenyl	piperidino	3,4-dichlorophenyl . 2HCl	259.6-260
78	(<i>R</i>)	4-methylphenyl	pyrrolidino	3,4-dichlorophenyl . 2HCl	249.6-250.1
79	(<i>RS</i>)	4-chlorophenyl	piperidino	3,4-dichlorophenyl . 2HCl	239.6-240.5
80	(<i>RS</i>)	4-methylphenyl	3-pyrrolino	3,4-dichlorophenyl . 2HCl	242-243

and are named as:

77. {1-(4-Methylbenzoyl)-2(*RS*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]}piperidine dihydrochloride salt.

IV. Representative compounds of Formula (I) where T and U = nitrogen; m and n = 1; $R^1=R^2=R^3$ = hydrogen; Q = $-\text{CH}_2-$; E = $-\text{NHC(O)NH}-$ and other groups are as defined below are:

[0053]



CPD #	Stereo- chem	Ar	F	R+R ⁴	R	R ⁴	Ar ¹	M.Pt.	Mass Spec. m/e
81	(R)	3-methoxyphenyl	bond		H	1-methylethyl	3,4-dichlorophenyl		478
82	(RS)	2,5-dimethoxyphenyl	bond		H	1-methylethyl	3,4-dichlorophenyl		508
83	(RS)	2-ethylphenyl	bond		H	1-methylethyl	3,4-dichlorophenyl		477
84	(RS)	3-methoxyphenyl	bond	cyclohexyl			3,4-dichlorophenyl		491.46
85	(RS)	3-ethylphenyl	bond		H	1-methylethyl	3,4-dichlorophenyl		476
86	(RS)	2-ethoxyphenyl	bond		H	1-methylethyl	3,4-dichlorophenyl		492
87	(RS)	2,3-dichlorophenyl	bond		H	1-methylethyl	3,4-dichlorophenyl		516
88	(RS)	3-bromophenyl	bond		H	1-methylethyl	3,4-dichlorophenyl		526
89	(RS)	phenyl	bond		H	1-methylethyl	4-chlorophenyl		414

CPD #	Stereo-chem	Ar	F	R+R ⁴	R	R ⁴	Ar ¹	M.Pt.	Mass Spec. m/e
90	(RS)	3-chloro-2-methyl-phenyl	bond		H	1-methylethyl	3,4-dichlorophenyl		496
91	(RS)	3-acetylphenyl	bond		H	1-methylethyl	3,4-dichlorophenyl		490
92	(RS)	phenyl	-CH ₂ -		H	1-methylethyl	3,4-dichlorophenyl		462
93	(R)	2,4,6-trimethylphenyl	bond		H	1-methylethyl	3,4-dichlorophenyl	271.1-275.5	
94	(RS)	4-methylthiophenyl	bond		H	2-methylpropyl	3,4-dichlorophenyl		508
95	(RS)	4-(2,2,2-trifluoroethyl) phenyl	bond		H	1-methylethyl	3,4-dichlorophenyl		532
96	(RS)	3-cyanophenyl	bond		H	2-methylpropyl	3,4-dichlorophenyl		487
97	(RS)	3-aminocarbonyl-phenyl	bond		H	1-methylethyl	3,4-dichlorophenyl . 2 HCl		492


and are named as:

82. 1-{1(*R*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-3-(3-methoxyphenyl)urea.
 84. 1-{2(*RS*)-[4-(3,4-Dichlorobenzyl)piperazin-1-yl]cyclohexyl}-3-(3-methoxyphenyl)urea.

92. I- $\{1(RS)$ -[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-3-benzylurea.

V. Representative compounds of Formula (I) where T = nitrogen; U = carbon; m and n =1; R=R¹=R²=R³=hydrogen; F = bond; and E = -NHC(O)NH- and other groups are as defined below are:

[0054]



CPD #	Stereo-chem	Ar	R ⁴	Q	Ar ¹	M.Pt.	Mass Spec.
98	(S)	3-methoxyphenyl	1-methylethyl	-CH ₂ -	3,4-dichlorophenyl . HCl	97.5-99.5	
99	(S)	3-methoxyphenyl	1,1-dimethylethyl	-CH ₂ -	3,4-dichlorophenyl	75.1-80.3	
100	(S)	4-methylphenyl	1,1-dimethylethyl	-CH ₂ -	3,4-dichlorophenyl . HCl	109.5-111	
101	(S)	3-carboxyphenyl	1,1-dimethylethyl	-CH ₂ -	3,4-dichlorophenyl		506
102	(S)	3-aminocarbonyl-phenyl	1,1-dimethylethyl	-CH ₂ -	3,4-dichlorophenyl		505
103	(S)	3,5-dimethoxy-phenyl	1-methylethyl	-CH ₂ -	3,4-dichlorophenyl . HCl		508
104	(RS)	3,4-dimethoxy-phenyl	1-methylethyl	-CH ₂ -	3,4-dichlorophenyl		508
105	(R)	3-methoxyphenyl	1-methylethyl	-CH ₂ -	3,4-dichlorophenyl		478
106	(S)	3-methoxyphenyl	1,1-dimethylethyl	-CH ₂ -	3,4-dichlorophenyl . TFA		492

5

10

15

20

25

30

35

40

45

50

CPD #	Stereo- chem	Ar	R ⁴	Q	Ar ¹	M.Pt.	Mass Spec.
					salt		


55 and are named as:

98. 1-{1(*S*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl}-3-(3-methoxyphenyl)urea.

EP 0 903 349 A2

VI. Representative compounds of Formula (I) where T and U = nitrogen; m and n =1; $R=R^1=R^2=R^3$ = hydrogen; Q = -CH₂-; E = -NHC(S)NH- and other groups are as defined below are:

[0055]



CPD #	Stereo-chem	Ar	F	R ⁴	Ar ¹	Mass Spec. m/e
107	(RS)	3-methylphenyl	bond	1-methylethyl	4-chlorophenyl	444
108	(RS)	2-fluorophenyl	bond	1-methylethyl	3,4-dichlorophenyl	482
109	(RS)	1-naphthyl	bond	1-methylethyl	4-chlorophenyl	480
110	(RS)	2-methoxyphenyl	bond	1-methylethyl	3,4-dichlorophenyl	494
111	(RS)	3-methylthiophenyl	bond	1-methylethyl	3,4-dichlorophenyl	510
112	(RS)	2,4-difluorophenyl	bond	2-methylpropyl	3,4-dichlorophenyl	514
113	(RS)	4-N,N-dimethylaminophenyl	bond	1-methylethyl	4-chlorophenyl	473
114	(RS)	4-methoxyphenyl	-CH ₂ -	1-methylethyl	3,4-dichlorophenyl	508
115	(RS)	4-trifluoromethylphenyl	bond	1-methylethyl	4-chlorophenyl	498
116	(RS)	3-chlorophenyl	bond	2-methylpropyl	3,4-dichlorophenyl	512


and are named as:

107. 1-{1(RS)-[4-(4-Chlorobenzyl)piperazin-1-yl]-2-methylpropyl}-3-(3-methylphenyl)-2-thiourea.

111. 1-{1(RS)-[4-(3,4-Dichlorobenzyl)piperazin-1-yl]-3-methylbutyl}-3-(2,4-difluorophenyl)-2-thiourea.

VII. Representative compounds of Formula (I) where T and U = nitrogen; m and n =1; $R=R^1=R^2=R^3$ = hydrogen; F = bond; Q = -CH₂-; E = -SO₂NH- and other groups are as defined below are:

[0056]



5

10

15

20

CPD #	Stereo-chem	Ar	R ⁴	Ar ¹	Mass Spec. m/e
117	(R)	4-fluorophenyl	1-methylethyl	3,4-dichlorophenyl	487
118	(RS)	4-chlorophenyl	1-methylethyl	4-chlorophenyl	469
119	(R)	4-nitrophenyl	1-methylethyl	3,4-dichlorophenyl	514
120	(R)	thiophen-2-yl	1-methylethyl	4-chlorophenyl	441
121	(RS)	4-trifluorophenyl	1-methylethyl	4-chlorophenyl	503
122	(RS)	2,4-dichlorophenyl	2-methylpropyl	3,4-dichlorophenyl	511
123	(RS)	3-bromophenyl	2-methylpropyl	3,4-dichlorophenyl	561
124	(R)	4-methylphenyl	1-methylethyl	4-chlorophenyl	449
125	(R)	2-chloro-4-fluorophenyl	1-methylethyl	4-chlorophenyl	487
126	(R)	2-nitro-4-trifluoromethyl-phenyl	1-methylethyl	4-chlorophenyl	480 (M ⁺ -CF ₃)
127	(RS)	3,4-dimethoxyphenyl	1-methylethyl	4-chlorophenyl	495

and are named as:

25

117. N-{1(*R*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-fluorobenzenesulfonamide.


120. N-{1(*RS*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-3-methylbutyl}-2,4-dichlorobenzenesulfonamide.

30

VIII. Representative compounds of Formula (I) where T and U = nitrogen; m and n = 1; R¹=R²=R³= hydrogen; F = bond; Q = -CH₂-; E = -NHC(O)- and other groups are as defined below are:

[0057]

35



40

45

50

55

CPD #	Stereo- chem	Ar	R+R ⁴	R ⁴	R	Ar ¹	M.Pt.	Mass Spec. m/e
128	(RS)	4-methylphenyl		methyl	H	3,4-dichlorophenyl . 2HCl	256.2-256.7	
129	(RS)	phenyl		methyl	H	3,4-dichlorophenyl . 2HCl	262.5-262.9	
130	1(R), 2(R)	4-methylphenyl	cyclopentyl			3,4-dichlorophenyl . 2HCl		446
131	1(R), 2(S)	4-methylphenyl	cyclopentyl			3,4-dichlorophenyl		

and are named as:


128. 2-(*RS*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-N-(4-methylphenyl)propionamide dihydrochloride salt.

130. 2-(*R*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-N-(4-methylphenyl)cyclopentane-1-(*R*)-carboxamide dihy-

drochloride salt.

IX. Representative compounds of Formula (I) where T = carbon; U = nitrogen; n = 1; R=R¹=R²=R³=hydrogen; F = bond; and Q = -CH₂-; E = -C(O)NH- and other groups are as defined below are:

[0058]




CPD #	Stereo- chem	Ar	E	R ⁴	m	R	Ar ¹	M.Pt.
132	(RS)	4-methoxyphenyl	-NHC(O)NH-	1-methyl/ethyl	0	H	3,4-dichlorophenyl	161.4-161.8
133	(RS)	4-methylphenyl	-C(O)NH-	1-methyl/ethyl	1	H	3,4-dichlorophenyl	233.9-235.5

and are named as:

133. N-{1-[1(*RS*)-[1-(3,4-Dichlorobenzyl)piperidin-4-ylmethyl]-2-methylpropyl]-4-methylbenzamide.

Certain compounds of Formula (I) are preferred.



15 [0059] A preferred group of compounds are those wherein:

n is 1;

m is 0 or 1;

F is a bond;

Q is an alkylene chain of between 1 to 6 carbon atoms inclusive, more preferably methylene; and

E is -C(O)N(R⁵)-, -SO₂N(R⁵)-, -N(R⁶)C(O)N(R⁵)- or -N(R⁶)C(O)-.

25 [0060] Within the above preferred group a more preferred group of compounds is that wherein:

R, R¹, R² and R³ are hydrogen; and

E is -C(O)N(R⁵)-, preferably -C(O)NH-. Another more preferred group of compounds is wherein:

R, R¹, R² and R³ are hydrogen; and

E is -N(R⁶)C(O)N(R⁵)-, preferably -NHC(O)NH-.

30 [0061] Within these preferred and more preferred groups of compounds a particularly preferred group of compounds is that wherein T and U are nitrogen;

Another particularly preferred group of compounds is that wherein T is nitrogen and U is carbon;

Yet another particularly preferred group of compounds is that wherein T is carbon and U is nitrogen;

[0062] Within the above preferred, more preferred and particularly preferred groups where T and U are both nitrogen or where T is nitrogen and U is carbon, most preferred compounds are those wherein:

40 R⁴ is alkyl or heteroalkyl, preferably 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl, more preferably 1-methylethyl or 1,1-dimethylethyl;

45 Ar is a heteroaryl or aryl ring, preferably a pyridin-2-yl, pyridin-3-yl, quinolin-3-yl or 5-methylthiophen-2-yl ring or a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH, more preferably a phenyl ring optionally substituted with one or two substituents selected from methyl, methoxy, fluoro, chloro, dimethylamino, acetyl, hydroxy, amino, methylenedioxy, -SO₂Me, 2-acetylaminoethyl, 2-[(R)-amino-3-methylbutyrylamino]ethyl, 2-aminoethyl, aminomethyl, hydroxymethyl, aminocarbonyl, -COOH, most preferably phenyl, 4-chlorophenyl, 3,4-difluorophenyl, 4-methylphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 4-hydroxyphenyl, 3,4-methylenedioxyphenyl, 4-methylsulfonylphenyl, 4-[(2-acetyl-amino)ethyl]phenyl, 4-[2-[(R)-amino-3-methylbutyrylamino]ethyl]phenyl, 4-(2-aminoethyl)phenyl, 4-(aminomethyl)phenyl, 4-(hydroxymethyl)phenyl, 3-aminocarbonylphenyl, 3-carboxyphenyl, 2,5-dimethoxyphenyl, 3,5-dimethoxyphenyl, 3,4-dimethoxyphenyl or 4-aminophenyl; and

50 Ar¹ is a heteroaryl or aryl ring, preferably 1-acetylbenzothien-3-yl, 3-methylbenzothien-2-yl or 5-nitrothiophen-3-yl, or a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro, or mono- or disubstituted amino, more preferably a phenyl ring substituted with one, or two substituents selected from methyl, methoxy, chloro, fluoro, trifluoromethyl or nitro, most preferably 4-nitrophenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 3,4-difluorophenyl, 2,3-dichlorophenyl, 3-methyl-4-nitrophenyl, 3-chloro-4-fluorophenyl or 3,4-dichlorophenyl.

[0063] Within the above preferred, more preferred and particularly preferred groups where T is carbon and U is nitrogen, most preferred compounds are those wherein:

R⁴ is alkyl or heteroalkyl, preferably methyl, 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl.

Ar is a heteroaryl or aryl ring, preferably a pyridin-2-yl, pyridin-3-yl, quinolin-3-yl or 5-methylthiophen-2-yl ring or a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH, more preferably a phenyl ring optionally substituted with one or two substituents selected from methyl, methoxy, fluoro, chloro, dimethylamino, acetyl, hydroxy, amino, methylenedioxy, -SO₂Me, 2-acetyl aminoethyl, 2-[(R)-amino-3-methylbutyrylamino]ethyl, 2-aminoethyl, aminomethyl, hydroxymethyl, aminocarbonyl or -COOH, most preferably phenyl, 4-chlorophenyl, 3,4-difluorophenyl, 4-methylphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 4-hydroxyphenyl, 3,4-methylenedioxyphenyl, 4-methylsulfonylphenyl, 4-[(2-acetyl amino)ethyl]phenyl, 4-{2-[(R)-amino-3-methylbutyrylamino]ethyl}phenyl, 4-(2-aminoethyl)phenyl, 4-(aminomethyl)phenyl, 4-(hydroxymethyl)phenyl, 3-aminocarbonyl-phenyl, 3-carboxyphenyl, 2,5-dimethoxyphenyl, 3,5-dimethoxyphenyl, 3,4-dimethoxyphenyl or 4-aminophenyl; and

Ar¹ is a heteroaryl or aryl ring, preferably 1-acetylindol-3-yl, 3-methylbenzothiophen-2-yl, 5-nitrothiophen-3-yl or a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro, or mono- or disubstituted amino, more preferably a phenyl ring substituted with one, or two substituents selected from methyl, methoxy, chloro, fluoro, trifluoromethyl or nitro, most preferably 4-nitrophenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 3,4-difluorophenyl, 2,3-dichlorophenyl, 3-methyl-4-nitrophenyl, 3-chloro-4-fluorophenyl or 3,4-dichlorophenyl.

[0064] Within the above preferred groups, particularly preferred compounds are those of formula I wherein n and m are 1; F is a bond; Q is an alkylene chain of between 1 to 6 carbon atoms inclusive; and E is -C(O)N(R⁵)-, -SO₂N(R⁵)-, -N(R⁵)C(O)N(R⁵)- or N(R⁵)C(O)-; particularly wherein R, R¹, R² and R³ are hydrogen; and E is -C(O)N(R⁵)-; particularly wherein T and U are both nitrogen; particularly wherein R⁴ is alkyl or heteroalkyl and R⁵ is hydrogen; particularly wherein Ar and Ar¹ are aryl; particularly wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl; particularly wherein Ar is a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH; and Ar¹ is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino; particularly wherein Ar is phenyl, 4-chlorophenyl, 3,4-difluorophenyl, 4-methylphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 4-hydroxyphenyl, 3,4-methylenedioxyphenyl, 4-methylsulfonylphenyl, 4-[(2-acetyl-amino)ethyl]phenyl, 4-{2-[(R)-amino-3-methylbutyrylamino]ethyl}phenyl, 4-(2-aminoethyl)phenyl, 4-(aminomethyl)-phenyl, 4-(hydroxymethyl)phenyl, 3-aminocarbonylphenyl, 3-carboxyphenyl, 2,5-dimethoxyphenyl, 3,5-dimethoxyphenyl, 3,4-dimethoxyphenyl or 4-aminophenyl; and Ar¹ is 4-nitrophenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 3,4-difluorophenyl, 2,3-dichlorophenyl, 3-methyl-4-nitrophenyl, 3-chloro-4-fluorophenyl or 3,4-dichlorophenyl; particularly wherein R⁴ is 1-methylethyl; Ar is 4-methylphenyl; Ar¹ is 3,4-dichlorophenyl; and Q is methylene; namely, N-{1-(S)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide dihydrochloride salt; or wherein R⁴ is 1,1-dimethylethyl; Ar is 4-methylphenyl; Ar¹ is 3,4-dichlorophenyl; and Q is methylene; namely, N-{1-(S)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethyl-propyl}-4-methylbenzamide dihydrochloride salt.

[0065] Among those preferred compounds of formula I in which n and m are 1; F is a bond; Q is an alkylene chain of between 1 to 6 carbon atoms inclusive; R, R¹, R² and R³ are hydrogen; E is -C(O)N(R⁵)-; T and U are both nitrogen; R⁴ is alkyl or heteroalkyl and R⁵ is hydrogen, such are preferred wherein Ar is an aryl ring; and Ar¹ is a heteroaryl ring; particularly wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl; particularly wherein Ar is a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH; and Ar¹ is 1-acetylindol-3-yl, 3-methylbenzothiophen-2-yl or 5-nitrothiophen-3-yl.

[0066] Also preferred among those preferred compounds of formula I in which n and m are 1; F is a bond; Q is an alkylene chain of between 1 to 6 carbon atoms inclusive; R, R¹, R² and R³ are hydrogen; and E is -C(O)N(R⁵)-; T and U are both nitrogen; R⁴ is alkyl or heteroalkyl and R⁵ is hydrogen, are those wherein Ar is a heteroaryl ring; and Ar¹ is an aryl ring; particularly wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl; particularly wherein Ar is pyridin-2-yl, pyridin-3-yl, quinolin-3-yl or 5-methylthiophen-2-yl; and Ar¹ is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino; particularly wherein Ar¹ is 4-nitrophenyl, 4-trifluoromethylphenyl, 4-

chlorophenyl, 3,4-difluorophenyl, 2,3-dichlorophenyl, 3-methyl-4-nitrophenyl, 3-chloro-4-fluorophenyl or 3,4-dichlorophenyl.

[0067] Among those preferred compounds of formula I, in which n and m are 1; F is a bond; Q is an alkylene chain of between 1 to 6 carbon atoms inclusive; R, R¹, R² and R³ are hydrogen; and E is -C(O)N(R⁵)-, such are preferred wherein T is nitrogen and U is carbon; particularly wherein R⁴ is alkyl or heteroalkyl and R⁵ is hydrogen; particularly wherein Ar and Ar¹ are aryl; particularly wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl; particularly wherein Ar is a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH; and Ar¹ is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino; particularly wherein Ar is phenyl, 4-chlorophenyl, 3,4-difluorophenyl, 4-methylphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 4-hydroxyphenyl, 3,4-methylenedioxyphenyl, 4-methylsulfonylphenyl, 4-[(2-acetyl-amino)ethyl]phenyl, 4-{2-[(R)-amino-3-methylbutylamino]ethyl}phenyl, 4-(2-aminoethyl)phenyl, 4-(aminomethyl)-phenyl, 4-(hydroxymethyl)phenyl, 3-aminocarbonylphenyl, 3-carboxyphenyl, 2,5-dimethoxyphenyl, 3,5-dimethoxyphenyl, 3,4-dimethoxyphenyl or 4-aminophenyl; and Ar¹ is 4-nitrophenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 3,4-difluorophenyl, 2,3-dichlorophenyl, 3-methyl-4-nitrophenyl, 3-chloro-4-fluoro-phenyl or 3,4-dichlorophenyl; particularly wherein R⁴ is 1-methylethyl; Ar is 4-methylphenyl; Ar¹ is 3,4-dichlorophenyl; and Q is methylene; and is named as N-{1-(S)-[4-(3,4-dichlorobenzyl)piperidin-1-ylmethyl]-2-methyl-propyl}-4-methylbenzamide dihydrochloride salt; or wherein R⁴ is 1-methylethyl; Ar is 4-(2-aminoethyl)phenyl; Ar¹ is 3,4-dichlorophenyl; and Q is methylene; and is named as, N-{1-(R)-[4-(3,4-dichlorobenzyl)piperidin-1-ylmethyl]-2-methyl-propyl}-4-(2-aminoethyl)benzamide dihydrochloride salt.

[0068] Among those preferred compounds of formula I in which n and m are 1; F is a bond; Q is an alkylene chain of between 1 to 6 carbon atoms inclusive; R, R¹, R² and R³ are hydrogen; and E is -C(O)N(R⁵)-; T is nitrogen and U is carbon; R⁴ is alkyl or heteroalkyl and R⁵ is hydrogen, such are preferred wherein Ar is a heteroaryl ring; and Ar¹ is an aryl; particularly wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl; particularly wherein Ar is pyridin-2-yl, pyridin-3-yl, quinolin-3-yl or 5-methylthiophen-2-yl; and Ar¹ is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino; particularly wherein Ar¹ is 4-nitrophenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 3,4-difluorophenyl, 2,3-dichlorophenyl, 3-methyl-4-nitrophenyl, 3-chloro-4-fluoro-phenyl or 3,4-dichlorophenyl; particularly wherein R⁴ is 1-methylethyl; Ar is 5-methylthiophen-2-yl; Ar¹ is 3,4-dichlorophenyl; and Q is methylene; and is named as, N-{1-(R)-[4-(3,4-dichlorobenzyl)piperidin-1-ylmethyl]-2-methyl-propyl}-5-methylthiophene-2-carboxamide hydrochloride salt.

[0069] Among those preferred compounds of formula I, in which n and m are 1; F is a bond; Q is an alkylene chain of between 1 to 6 carbon atoms inclusive; R, R¹, R² and R³ are hydrogen; and E is -C(O)N(R⁵)-, such are preferred wherein T is carbon and U is nitrogen; particularly wherein R⁴ is alkyl or heteroalkyl; and R⁵ is hydrogen; particularly wherein Ar is an aryl or heteroaryl ring; and Ar¹ is an aryl ring; particularly wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl; particularly wherein Ar is a pyridin-2-yl, pyridin-3-yl, quinolin-3-yl or 5-methylthiophen-2-yl ring, or a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH; and Ar¹ is a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino.

[0070] Among those preferred compounds of formula I in which n and m are 1; F is a bond; Q is an alkylene chain of between 1 to 6 carbon atoms inclusive; and E is -C(O)N(R⁵)-, -SO₂N(R⁵)-, -N(R⁶)C(O)N(R⁵)- or N(R⁶)C(O)-, such are preferred wherein R, R¹, R² and R³ are hydrogen; and E is -N(R⁶)C(O)N(R⁵)-; particularly wherein T and U are both nitrogen; particularly wherein R⁴ is alkyl or heteroalkyl; and R⁵ and R⁶ are hydrogen; particularly wherein Ar and Ar¹ are aryl; particularly wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl, particularly wherein Ar is a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH; and Ar¹ is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino; particularly wherein Ar is phenyl, 4-chlorophenyl, 3,4-difluorophenyl, 4-methylphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 4-hydroxyphenyl, 3,4-methylenedioxyphenyl, 4-methylsulfonylphenyl, 4-[(2-acetyl-amino)ethyl]phenyl, 4-{2-[(R)-amino-3-methylbutylamino]ethyl}phenyl, 4-(2-aminoethyl)phenyl, 4-(aminomethyl)-phenyl, 4-(hydroxymethyl)phenyl, 3-aminocarbonylphenyl, 3-carboxyphenyl, 2,5-dimethoxyphenyl, 3,5-dimethoxyphenyl, 3,4-dimethoxyphenyl or 4-aminophenyl; and Ar¹ is 4-nitrophenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 3,4-difluorophenyl, 2,3-dichlorophenyl, 3-methyl-4-nitrophenyl, 3-chloro-4-fluoro-phenyl or 3,4-dichlorophenyl; particularly wherein R⁴ is 1-methylethyl; Ar is 3-methoxyphenyl; Ar¹ is 3,4-dichlorophenyl;

and Q is methylene; namely, 1-{1-(R)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-3-(3-methoxyphenyl)urea.

[0071] Among those preferred compounds of formula I in which n and m are 1; F is a bond; Q is an alkylene chain of between 1 to 6 carbon atoms inclusive; R, R¹, R² and R³ are hydrogen; and E is -N(R⁶)C(O)N(R⁵)-, such are preferred wherein T is nitrogen and U is carbon; particularly wherein R⁴ is alkyl or heteroalkyl; and R⁵ and R⁶ are hydrogen; particularly wherein Ar and Ar¹ are aryl; particularly wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl; particularly wherein Ar is a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH; and Ar¹ is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino; particularly wherein Ar is phenyl, 4-chlorophenyl, 3,4-difluorophenyl, 4-methylphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 4-hydroxyphenyl, 3,4-methylenedioxyphenyl, 4-methylsulfonylphenyl, 4-[(2-acetyl-amino)ethyl]phenyl, 4-[2-[(R)-amino-3-methylbutyrylamino]ethyl]phenyl, 4-(2-aminoethyl)phenyl, 4-(aminomethyl)-phenyl, 4-(hydroxymethyl)phenyl, 3-aminocarbonylphenyl, 3-carboxyphenyl, 2,5-dimethoxyphenyl, 3,5-dimethoxyphenyl, 3,4-dimethoxyphenyl or 4-aminophenyl; and Ar¹ is 4-nitrophenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 3,4-difluorophenyl, 2,3-dichlorophenyl, 3-methyl-4-nitrophenyl, 3-chloro-4-fluoro-phenyl or 3,4-dichlorophenyl; particularly wherein R⁴ is 1-methylethyl; Ar is 3-methoxyphenyl; Ar¹ is 3,4-dichlorophenyl; and Q is methylene; namely, 1-{1-(R)-[4-(3,4-dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl}-3-(3-methoxyphenyl)urea.

[0072] Among those preferred compounds of formula I in which n and m are 1; F is a bond; Q is an alkylene chain of between 1 to 6 carbon atoms inclusive; R, R¹, R² and R³ are hydrogen; and E is -N(R⁶)C(O)N(R⁵)-, such are preferred wherein T is carbon and U is nitrogen; particularly wherein R⁴ is alkyl or heteroalkyl; and R⁵ and R⁶ are hydrogen; particularly wherein Ar and Ar¹ are aryl; particularly wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl; particularly wherein Ar is a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH; and Ar¹ is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino.

[0073] Finally, preferred compounds of formula are such wherein T is carbon and U is nitrogen; n is 1; m is 0; F is a bond; Q is an alkylene chain; and E is -C(O)N(R⁵)-, -SO₂N(R⁵)-, -N(R⁶)C(O)N(R⁵)- or N(R⁶)C(O)-; particularly wherein R, R¹, R² and R³ are hydrogen; and E is -C(O)N(R⁵)-; particularly wherein R⁴ is alkyl or heteroalkyl; and R⁵ is hydrogen; particularly wherein Ar is an aryl or heteroaryl ring; and Ar¹ is an aryl ring; particularly wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl; particularly wherein Ar is pyridin-2-yl, pyridin-3-yl, quinolin-3-yl or 5-methylthiophen-2-yl, or a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH; and Ar¹ is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino.

[0074] Also, among those preferred compounds of formula I in which T is carbon and U is nitrogen; n is 1; m is 0; F is a bond; Q is an alkylene chain; and E is -C(O)N(R⁵)-, -SO₂N(R⁵)-, -N(R⁶)C(O)N(R⁵)- or N(R⁶)C(O)-, such are preferred wherein R, R¹, R² and R³ are hydrogen; and E is -(NR⁶)C(O)N(R⁵)-; particularly wherein R⁴ is alkyl or heteroalkyl; and R⁵ and R⁶ are hydrogen; particularly wherein Ar is an aryl or heteroaryl ring; and Ar¹ is an aryl ring; particularly wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl; particularly wherein Ar is pyridin-2-yl, pyridin-3-yl, quinolin-3-yl or 5-methylthiophen-2-yl, or a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH; and Ar¹ is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino.

[0075] Exemplary particularly preferred compounds of the invention at present are:

N-{1-(S)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide dihydrochloride salt.

N-{1-(S)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-4-methylbenzamide dihydrochloride salt.

1-{1-(R)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-3-(3-methoxyphenyl)urea.

N-{1-(RS)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-3,4-methylenedioxybenzamide.

N-{1-(S)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-4-methylsulfonylbenzamide dihydrochloride salt.

N-{1-(S)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-4-acetylbenzamide dihydrochloride salt.

N-{1-(*S*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-4-dimethylaminobenzamide dihydrochloride salt.

N-{1-(*S*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-5-methylthiophene-2-carboxamide dihydrochloride salt.

N-{1-(*RS*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-4-methoxybenzamide.

N-{1-(*RS*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-3-cyanobenzamide.

N-{1-(*RS*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-3,4-difluorobenzamide.

N-{1-(*RS*)-[3-Methyl-4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide dihydrochloride salt.

N-{4-(*R*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl}-1-[4-(2-acetylaminoethyl)]benzamide dihydrochloride salt.

4-[2-(2-(*R*)-Amino-3-methylbutyrylamino)ethyl]-N-{1-[4-(3,4-dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl}benzamide dihydrochloride salt.

N-{4-(*R*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2,2-dimethylpropyl}-1-[4-(2-aminoethyl)]benzamide dihydrochloride salt.

N-{4-(*R*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2,2-dimethylpropyl}-1-(4-aminomethyl)benzamide dihydrochloride salt.

1-{1-(*RS*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-3-(3-aminocarbonylphenyl)urea.

N-{4-(*R*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2,2-dimethylpropyl}-1-quinoline-3-carboxamide hydrochloride salt.

1-{1-(*S*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2,2-dimethylpropyl}-3-(3-carboxyphenyl)urea.

1-{1-(*S*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2,2-dimethylpropyl}-3-(3-aminocarbonylphenyl)urea.

1-{1-(*S*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl}-3-(3,5-dimethoxyphenyl)urea hydrochloride salt.

1-{1-(*RS*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl}-3-(3,4-dimethoxyphenyl)urea.

N-{1-(*RS*)-[4-(4-Chlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide.

N-{1-(*RS*)-[4-(4-Nitrobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide.

N-{1-(*RS*)-[4-(3,4-Difluorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide.

N-{1-(*RS*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-4-methylbenzamide.

N-{1-(*RS*)-[4-(2,3-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide.

N-{1-(*RS*)-[4-(3-Methyl-4-nitrobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide.

N-{1-(*RS*)-[4-(3-Chloro-4-fluorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide.

N-{1-(*R*)-[4-(3-Chloro-4-fluorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-4-methylbenzamide.

N-{1-(*R*)-[4-(3-Methylbenzothiophen-2-ylmethyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-4-methylbenzamide.

N-{1-(*R*)-[4-(1-Acetylindol-3-ylmethyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-4-methylbenzamide.

N-{1-(*R*)-[4-(5-Nitrothiophen-3-ylmethyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-4-methylbenzamide.

N-{1-(*R*)-[4-(4-Nitrobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide.

N-{4-(*R*)-[4-(4-Nitrobenzyl)piperidin-1-ylmethyl]-2,2-dimethylpropyl}-1-pyridine-2-carboxamide hydrochloride salt.

N-{4-(*R*)-[4-(4-Nitrobenzyl)piperidin-1-ylmethyl]-2,2-dimethylpropyl}-1-pyridine-2-carboxamide.

N-{4-(*R*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-1-quinoline-3-carboxamide.

1-{1-(*R*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl}-3-(3-methoxyphenyl)urea.

N-{4-(*R*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl}-1-(3-methyl)thiophene-2-carboxamide hydrochloride salt.

N-{4-(*R*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl}-1-[4-(2-aminoethyl)]benzamide dihydrochloride salt.

N-{4-(*RS*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl}-1-(4-methyl)benzamide.

N-{4-(*R*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2,2-dimethylpropyl}-1-(4-methyl)benzamide hydrochloride salt.

N-{4-(*R*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2,2-dimethylpropyl}-1-(4-methylsulfonyl)benzamide.

N-{4-(*R*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2,2-dimethylpropyl}-1-(5-methylthiophene-2-carboxamide hydrochloride salt.

N-{4-(*R*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2,2-dimethylpropyl}-1-(4-hydroxymethyl)benzamide.

1-{1-(*R*)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2,2-dimethylpropyl}-3-(3-methoxyphenyl)urea trifluoroacetate salt.

[0076] The compounds of the present invention can be prepared in a number of ways known to one skilled in the art. Preferred methods include, but are not limited to, the general synthetic procedures described below.

[0077] The starting materials and reagents used in preparing these compounds are either available from commercial

suppliers such as Aldrich Chemical Co., (Milwaukee, Wisconsin, USA), Bachem (Torrance, California, USA), Emka-Chemie, or Sigma (St. Louis, Missouri, USA) or are prepared by methods known to those skilled in the art following procedures set forth in references such as *Fieser and Fieser's Reagents for Organic Synthesis*, Volumes 1-17 (John Wiley and Sons, 1991); *Rodd's Chemistry of Carbon Compounds*, Volumes 1-5 and Supplementals (Elsevier Science Publishers, 1989), *Organic Reactions*, Volumes 1-40 (John Wiley and Sons, 1991), *March's Advanced Organic Chemistry*, (John Wiley and Sons, 1992), and *Larock's Comprehensive Organic Transformations* (VCH Publishers Inc., 1989). These schemes are merely illustrative of some methods by which the compounds of this invention can be synthesized, and various modifications to these schemes can be made and will be suggested to one skilled in the art having referred to this disclosure.

[0078] The starting materials and the intermediates of the reaction may be isolated and purified if desired using conventional techniques, including but not limited to filtration, distillation, crystallization, chromatography, and the like. Such materials may be characterized using conventional means, including physical constants and spectral data.

Synthesis of Compounds of Formula (I)

[0079] In general, compounds of Formula (I) where n , m , R , R^1 , R^2 , R^3 , R^4 , R^5 , Q and Ar^1 are as defined above are prepared from aminoalkyl derivatives of Formulae II(a-c) and carboxyalkyl derivatives of Formulae II(d-f) as shown in Fig. 1 below.




Fig. 1


[0080] Synthesis of compounds of Formulae II(a-f) and their conversion to compounds of Formula (I) are described in detail in Schemes A-E and F-J, respectively.

Synthesis of compounds of Formulae II(a-f)

Preparation of Compounds of Formula IIa

[0081] A compound of Formula IIa where n is 1 or 2, m is at least 1 and R , R^1 , R^2 , R^3 , R^4 , R^5 , Q and Ar^1 are as defined above is prepared as shown in Scheme A below.

Scheme A




[0082] In general, compounds of Formula IIa are prepared in two steps by first converting a compound of formula 1 or 2 to an N-protected aminoalkyl derivative of formula 3 via methods (a) or (b) respectively, followed by removal of the amino protecting group in 3, as described in detail below.

Preparation of compounds of formula 3

Method (a)


[0083] In method (a), an N-protected aminoalkyl derivative of formula 3 where PG is an amino protecting group (e.g., *tert*-butoxycarbonyl (BOC), benzyloxycarbonyl (CBZ), benzyl, and the like) is prepared by reacting a compound of formula 1 with a compound of formula 4



where X is an aldehyde (-CHO), ketone (-C(O)R where R is alkyl), carboxy (-COOH) or a reactive carboxy derivative e.g., acid halide. The reaction conditions employed depend on the nature of the X group. If X is an aldehyde or a ketone group, the reaction is carried out under reductive amination reaction conditions i.e., in the presence of a suitable reducing agent (e.g., sodium cyanoborohydride, sodium triacetoxyborohydride, and the like) and an organic acid (e.g., glacial acetic acid, trifluoroacetic acid, and the like) at ambient temperature to give 3 directly. Suitable solvents for the reaction are halogenated hydrocarbons (e.g., 1,2-dichloroethane, chloroform, and the like). If X is a carboxy group, the reaction is carried out in the presence of a suitable coupling agent (e.g., N,N-dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide, and the like) in a suitable organic solvent (e.g., methylene chloride, tetrahydrofuran, and the like) to give an amide intermediate. Reduction of the amide intermediate with a suitable reducing agent (e.g., diborane, lithium aluminum hydride, and the like) in an ethereal organic solvent such as ether or tetrahydrofuran then provides a compound of formula 3. If X is an acid derivative such as an acid chloride, the reaction is carried out in the presence of a suitable base such as triethylamine, pyridine in an organic solvent (e.g., methylene chloride, dichloroethane, N,N-dimethylformamide, and the like) to give an amide intermediate which is reduced to the corresponding compound of formula 3 as described above.

[0084] In general, compounds of formula 4 are either commercially available or they can be prepared by methods well known in the field of organic chemistry. Some examples of such procedures are illustrated and described in detail

below.



(i) An aldehyde of 4 (X is a -CHO) where m-1 = 0 is conveniently prepared from the corresponding natural or unnatural α -amino acid of formula 4 where m-1 = 0 and X is a carboxy group by reduction of the carboxy group to an aldehyde group with a suitable reducing agent such as DIBAL-H[®]. An aldehyde of formula 4 where m-1 = 1 or 2 can be prepared, if desired, from an aldehyde or ketone (X = -COR where R is alkyl) of formula 4 where m-1 = 0 under Wittig reaction conditions. For example, an aldehyde 4 where m-1 = 1 and R is hydrogen or alkyl is prepared by condensing the corresponding aldehyde or ketone of formula 4 where m-1 = 0 with a Wittig reagent derived from chloromethyl methyl ether, followed by acidic hydrolysis of the resulting enol ether intermediate. An aldehyde 4 where m-1 = 2 and R are hydrogen or alkyl can be prepared by condensing the corresponding aldehyde or ketone 4 where m-1 = 0 with a Wittig reagent derived from bromoacetate or 2-bromopropionate respectively, followed by sequential reduction of the double bond and the ester group in the resulting α,β -unsaturated ester intermediate. The double bond is reduced under hydrogenation reaction conditions and the ester group is reduced to the aldehyde group with reducing agent such as DIBAL-H[®]. The ketone of formula 4 where m-1 = 0 can be prepared from the α -amino acids of formula 4 by converting the α -amino acids 4 to a Weinreb amide, followed by treatment with an organometallic reagent such as a Grignard reagent or an organolithium reagent of formula RMgBr or RLi (where R is an alkyl group) respectively.

Generally, both natural and unnatural amino acids are commercially available from vendors such as Aldrich

and Bachem. Examples of unnatural amino acids include, homoserine, homocysteine, N- α -methylarginine, norleucine, N-methylisoleucine, phenylglycine, hydroxyproline, pyroglutamine, omithine, 2-aminoisobutyric acid, 2-aminobutyric acid, β -cyclohexylalanine, 3-(1-naphthyl)alanine, 3-(2-naphthyl)alanine, citrulline, pipecolic acid, piperazic acid, 4-chlorophenylalanine, 4-fluorophenylalanine and sarcosine. Synthesis of α -amino acids 4 where R³ and R⁵ together form a morpholino ring and thiomorpholino ring and R⁴ is hydrogen is described in Kogami, Y., Okawa, K., *Bull. Chem. Soc. Jpn.*, 60, 2963, (1987).

(ii) Compounds of formula 4 where X is a carboxy group and m-1>0 can be prepared from the corresponding aldehydes of formula 4 (X is -CHO), prepared as described in (i) above, by oxidation of the aldehyde group with a suitable oxidizing agent (e.g., potassium permanganate and the like). Alternatively, they can be prepared from the α,β -unsaturated esters formed in the Wittig reaction, see (i) above, by reduction of the double bond followed by the hydrolysis of the ester group by methods well known in the art.

(iii) Compounds of formula 4 where X is -C(O)R (where R is alkyl) and m-1 = 0, 1 or 2 can be prepared by alkylating the corresponding aldehyde of formula 4 (X is -CHO) with a Grignard reagent followed by oxidation of the resulting alcohol with a suitable oxidizing agent such as pyridinium dichromate and the like. Alternatively, they can be prepared from the corresponding acid of formula 4 as described in (i) above.

(iv) Compounds of formula 4 where X is an acid derivative e.g., an acid chloride, can be prepared from the corresponding acids of formula 4 (X is -COOH), prepared as described in (iii) above, by chlorinating the carboxy group with a suitable chlorinating agent (e.g., oxalyl chloride, thionyl chloride and the like) in a suitable organic solvent such as methylene chloride and the like.

[0085] Alternatively, a compound of formula 3 can be prepared directly by reacting a compound of formula 1 with an alkylating agent of formula 5



5


where Y is a leaving group under alkylating conditions such as halo (e.g., chloro, bromo or iodo) or sulfonyloxy group (e.g., methylsulfonyloxy or 4-methylphenylsulfonyloxy or trifluoromethylsulfonyloxy). The reaction is carried out in the presence of a base such as sodium carbonate, sodium hydride, triethylamine and the like. Suitable solvents are aprotic organic solvents such as tetrahydrofuran, N,N-dimethylformamide, and the like.

[0086] In general, compounds of formula 5 where Y is a halo or a sulfonyloxy group can be prepared from compounds of formula 4 by reducing the aldehyde, ketone or carboxy group to an alcohol, followed by treatment with a suitable halogenating agent (e.g., thionyl chloride, thionyl bromide, carbon tetrabromide in the presence of triphenylphosphine, and the like) or sulfonylating agent (e.g., methylsulfonyl chloride, para-toluenesulfonyl chloride and triflic anhydride) respectively. Suitable aldehyde, ketone or carboxy reducing agents include lithium aluminum hydride, borane and the like.

[0087] In some instances, a compound of Formula IIa can be prepared by reacting a compound of formula 1 with a conjugated nitro-olefin under Michael addition reaction conditions, followed by reduction of the nitro group under standard hydrogenation reaction conditions. Conjugated nitro-olefins are commercially available or can be prepared by known literature procedures e.g., see Corey, E.J. et al., *J. Am. Chem. Soc.*, 100(19), 8294-5, (1978). A detailed description of the synthesis of an N-alkylaminopiperazine of Formula IIa by this method is given in Example 2.

Method (b)

[0088] In method (b), an N-protected aminoalkyl derivative of formula 3 is prepared by reacting a compound of formula 2 with a compound of formula 6



6

(where J is an X or Y group as defined above) utilizing the reaction conditions described in method (a) above. Method (b) is particularly suitable for preparing compounds of Formula IIa where Q contains an amido or a carbonyl group.

[0089] In general, compounds of formula 6 are commercially available or can be prepared by methods well known in the art. For example, aralkyl halides and aralkyl acids such as benzyl bromide, 3,4-dichlorobenzyl bromide, phenylacetic acids and 2-phenylpropionic acids are commercially available. Others can be prepared from suitable starting mate-


rials such as phenylacetic acid, phenylpropanol, 2-pyridineethanol, nicotinic acid etc., by following procedures described for the synthesis of compounds of formula 4 and 5 in method (a) above. Compounds of formula 6 where Q is an alkylene chain interrupted by an amido group and J is a halo or sufonyloxy group can be prepared by following the procedures described in U.S. Pat. No. 4,880,808.

Conversion of compounds of formula 3 to compounds of Formula IIa

[0090] In Step 2, the N-protected aminoalkyl derivatives 3, formed in Step 1 via method (a) or (b), are converted to a compound of Formula IIa by removal of the amino protecting group. The conditions utilized depend on the nature of the protecting group. For example, if the protecting group is the *tert*-butoxycarbonyl group it is removed under acidic hydrolysis reaction condition whereas if it is the benzyl group it is removed under catalytic hydrogenation reaction conditions.

[0091] A compound of Formula IIa where R^5 is other than hydrogen can be prepared, if desired, by alkylating the corresponding compound of Formula IIa where R^5 is hydrogen with an alkylating agent R^5Y where Y is a leaving group under alkylating conditions, utilizing the reaction conditions described in method (a) of Scheme A.

[0092] Compounds of formula 1 and 2 are prepared as shown below by reacting a piperazine or homopiperazine of formula 7 with a compound of formula 6, or 4 or 5 respectively, followed by the removal of the amino protecting group, utilizing the reaction conditions described in method (a) above.




[0093] Piperazines and homopiperazines of formula 7 such as piperazine, 2 or 3-methylpiperazines, homopiperazine are commercially available. Piperazines 7 can also be prepared by following the procedures described in the European Pat. Pub. No. 0,068,544 and U.S. Pat. No. 3,267,104. Detailed descriptions of the synthesis of a compound of formula 1 where $n = 1$ by this method is given in Examples 1, 5 and 7.

Preparation of Compounds of Formula IIb

[0094] A compound of Formula IIb where m is at least 1 and n , R , R^1 , R^2 , R^3 , R^4 , R^5 , Q and Ar^1 are as defined in the Summary of the invention can be prepared from a compound of formula 8, where n is 0, 1 or 2 respectively, as illustrated in Scheme B below.

Scheme B



[0095] In general, an aminoalkyl derivative of Formula IIb is prepared by reacting a compound of formula 8 with a compound of formula 4 or 5 (see Scheme A) to give an N-protected aminoalkyl derivative of formula 9, followed by removal of the amino protecting group. The conversion of a compound of formula 8 to a compound of Formula IIb is carried out under the reaction conditions described in method (a) of Scheme A above.

[0096] Compounds of formula 8 where n is 0, 1 or 2 can be prepared from suitably N-protected pyrrolidinones, piperidinones or 4-keto-octahydroazepines respectively, by known procedures. Some examples of such procedures are described below:

(i) Compounds of formula 8 where n is 0 or 1 and Q is an alkylene chain are prepared by reacting a suitably N-protected-3-pyrrolidinone or N-protected-4-piperidinone respectively, with a Wittig reagent $Br(Ph)_3P^+-alkylene-Ar^1$ to give an alkene intermediate. Reduction of the olefinic bond, followed by removal of the N-protecting group then provides compounds of formula 8.


[0097] 4-Hydroxypiperidines, 3-pyrrolidinols, 3-pyrrolidinones and 4-piperidinones are commercially available. 4-keto-octahydroazepine can be prepared from 2,4-diketo-N-benzylhexahydroazepine ((see Hong Hu G. and Erik Jagdmann Jr., *Tet. Lett.*, 36(21), 3659-62 (1995)) by known procedures.

[0098] Detailed descriptions of the synthesis of compounds of Formula IIb by this method are given in Examples 3 and 4.

Preparation of Compounds of Formula IIc

[0099] A compound of Formula IIc where m is 0 or 1, R^3 is hydrogen and n , R , R^1 , R^2 , R^4 , R^5 , Q and Ar^1 are as defined in the Summary of the invention can be prepared from a compound of formula 14 or 10 respectively, as illustrated in Scheme C below.

Scheme C



[0100] A compound of Formula IIc where m is 1 can be prepared, as shown in method (a), by reacting a compound of formula 10 with a phosphonate ylide of formula 11 under Wittig reaction conditions, i.e., in the presence of a strong non-nucleophilic base (e.g., sodium hydride, sodium amide, and the like) and in a suitable aprotic organic solvent (e.g., tetrahydrofuran and the like) to give an α,β -unsaturated ester of formula 12. The α,β -unsaturated ester 12 is converted to the corresponding alcohol derivative 13a (m = 1) by first converting 12 to an aldehyde, followed by treatment with an organometallic reagent such as a Grignard reagent or an organolithium reagent of formula R^4MgBr or R^4Li , respectively. The double bond is reduced under hydrogenation reaction conditions and the ester group is reduced to the aldehyde group with a suitable reducing agent such as DIBAL- H° . The alcohol 13a is then converted to a compound of Formula IIc by oxidation of the alcohol group to the ketone group, followed by treatment with an amine of formula $NH(R^5)$ under reductive amination reaction conditions. The oxidation reaction is carried in with a suitable oxidizing reagents such as pyridinium dichromate in an aprotic solvent such as dimethylformamide and the like.

[0101] A compound of Formula IIc where m is 0 can be prepared, as shown in method (b) from a compound of formula 14, by converting 14 to the corresponding alcohol derivative 13b (m = 0) by reduction of the ester group to the aldehyde followed by treatment with a suitable organometallic reagent. Compound 13b is then converted to a compound of IIc where m is 0 by carrying out the oxidation and reductive amination steps, utilizing the reaction conditions described above. Compounds of Formula IIc where m is 0 can also be prepared by the procedures described in PCT application Publication No. WO 92/12128.

[0102] Compounds of formula 10 where n is 0, 1 or 2 are prepared by N-alkylating a 3-pyrrolidone, a 4-piperidone or a 4-keto-octahydroazepine respectively, with a compound of formula Ar^1-Q-Y where Y is a leaving group under alkylating conditions as described in method (a) (2) of Scheme A.

[0103] Compounds of formula 14 (n = 1) are prepared by N-alkylating an ethyl isonipecotate with a compound of for-


mula $\text{Ar}^1\text{-Q-Y}$ where Y is a leaving group under alkylating conditions as described in method (a) of Scheme A.

[0104] Detailed descriptions of the synthesis of a compound of Formula IIc where m is 0 or 1 is given in Examples 9 and 10 respectively.

5 Preparation of Compounds of Formula II d and II e

[0105] A carboxyalkyl derivative of Formula II d (U = N) and II e (U = C) where m, n, R^1 , R^2 , R^3 , R^4 , Q and Ar^1 are as defined in the Summary of the invention can be prepared from a compound of formula 1 or 8 respectively, as illustrated in Scheme D below.

Scheme D




[0106] A carboxy derivative of Formula II d or II e is prepared, as shown above, by reacting a compound of formula 1 or 8 with an alkylating agent of formula 15 where Y is halo or sulfonyloxy group, followed by hydrolysis of the ester group. The alkylation reaction is carried out under the reaction conditions described previously (see scheme A method (a)). The hydrolysis of the ester group is carried out in the presence of an aqueous base (e.g., sodium hydroxide, lithium hydroxide, and the like) in an alcoholic organic solvent such as methanol, ethanol, and the like. The reaction proceeds either at ambient temperature or upon heating. Alternatively, a carboxyethyl derivative of Formula II d or II e where R^3 is hydrogen is prepared by reacting a compound of formula 1 or 8 with an α,β -unsaturated ester of formula 16 under Michael addition reaction conditions i.e., in the presence of a suitable base such as methoxide and in a protic organic solvent (e.g., methanol, ethanol and the like) to give a 3-propionate derivative of formula 17a or 17b, respectively. Hydrolysis of the ester group in 17a or 17b then provides the corresponding carboxyethyl derivative of Formula II d or II e respectively, where R^3 is hydrogen.

[0107] Compounds of formula 1 or 8 are prepared as described previously in Schemes A and B respectively. Compounds of formula 15 and 16 are either commercially available or can be prepared by methods known in the art. For example, halo acids and α,β -unsaturated ester such as methyl 2-bromo-2-methylpropionate, methyl 2-bromopropionate, methyl 3-bromo-2-methylpropionate, methyl α -bromophenylacetate, methyl methacrylate are commercially available. A detailed description of the synthesis of a carboxyethylpiperazine of Formula II d by this method is given in Example 6.

Preparation of Compounds of Formula II f

[0108] A carboxyalkyl derivative of Formula II f where n, m, R^1 , R^2 , R^3 , R^4 , Q and Ar^1 are as defined in the Summary of the invention can be prepared from a compound of formula 17 or 18 respectively, as illustrated in Scheme E below.

Scheme E



[0109] A carboxyalkyl derivative of Formula II(f) can be prepared by reacting a compound of formula 10 or 18 with a Wittig reagent of formula $\text{Br}-(\text{Ph}_3\text{P})(\text{CHR})_m\text{CR}^3\text{R}^4\text{CO}_2\text{Et}$, followed by reduction of the double bond and hydrolysis of the ester group to the acid in the resulting unsaturated ester 19, as described previously. Alternatively, compounds of Formula II(f) can be prepared from 18 (where R is hydrogen or alkyl) by following the reaction procedures described for the synthesis of compound 4 where X is carboxy in Scheme A, method (a).

[0110] Compounds of formula 18 can be prepared from compounds of formula 14 by methods well known in the art.


[0111] Compounds of Formula II(d-f) are used for the synthesis of compound of Formula (I) where E is an inverse amide i.e., $-\text{N}(\text{R}^6)\text{CO}-$.

40 Synthesis of Compounds of Formula (I) from Compounds of Formulae II(a-f)

[0112] Compounds of Formula (I) are prepared from compounds of Formulae II(a-f) as described in Schemes F - J below.

[0113] Compounds of Formula (I) where E is $-\text{C}(\text{O})\text{N}(\text{R}^5)-$ are prepared as described in Scheme F below:

Scheme F




[0114] A compound of Formula (I) where E is an amide group can be prepared, either:

(i) by reacting a compound of Formula II(a-c) with an acylating reagent Ar-F-C(O)L , where L is a leaving group under acylating conditions, such as a halo (particularly Cl or Br) or imidazolidine. Suitable solvents for the reaction include aprotic polar solvents (e.g., dichloromethane, THF, dioxane and the like). When an acyl halide is used as the acylating agent the reaction is carried out in the presence of a non-nucleophilic organic base (e.g., triethylamine or pyridine, preferably pyridine); or

(ii) by heating a compound of Formula II (a-c) with an acid anhydride. Suitable solvents for the reaction are tetrahydrofuran, dioxane and the like. Detailed descriptions of the conversion of a compound of Formula IIa and IIb to compounds of Formula (I) where n is 1 and E is $-\text{C(O)NH}-$ is given in Examples 1, 3, 4, 5 and 10.

[0115] Compounds of Formula (I) where E is $-\text{N(R}^6\text{)C(O)N(R}^5\text{)}-$ or $-\text{N(R}^6\text{)C(S)N(R}^5\text{)}-$ are prepared as described in Scheme G below:

Scheme G



[0116] A compound of Formula (I) where E is a urea/thiourea group can be prepared, either:

(i) by reacting a compound of Formula II(a-c) with an activating agent such as carbonyl diimidazole/ thiocarbonyl diimidazole, followed by nucleophilic displacement of the imidazole group with a primary or secondary amine. The reaction occurs at ambient temperature. Suitable solvents include polar organic solvents (e.g., tetrahydrofuran, dioxane and the like);


(ii) by reacting a compound of Formula II (a-c) with a carbamoyl/thiocarbamoyl halide. The reaction is carried out in the presence of a non-nucleophilic organic base. Suitable solvents for the reaction are dichloromethane, 1,2-dichloroethane, tetrahydrofuran or pyridine; or

(iii) by reacting a compound of formula II (a-c) with an isocyanate/isothiocyanate in an aprotic organic solvent (e.g., benzene, tetrahydrofuran, dimethylformamide and the like).

[0117] Detailed descriptions of the conversion of a compound of Formula IIa to a compound of Formula (I) where n is 1 and E is $-\text{NHC(O)NH}-$ or $-\text{N(R}^6\text{)C(S)N(R}^5\text{)}-$ is given in Examples 2 and 8. A detailed description of the conversion of a compound of Formula IIc where m is 0 to a compound of Formula (I) where E is $-\text{NHC(O)NH}-$ is given in Example 9.

[0118] Compounds of Formula (I) where E is $-\text{SO}_2\text{N(R}^5\text{)}-$ are prepared as described in Scheme H below:

Scheme H




[0119] A compound of Formula (I) where E is a sulfonamido group can be prepared by reacting a compound of For-

mula II(a-c) with a sulfonyl halide, utilizing the reaction conditions described in method (i) of Scheme F. Sulfonyl halides are commercially available or may be prepared by methods such as those described in (1) Langer, R. F.; *Can. J. Chem.* 61, 1583-1592, (1983); (2) Aveta, R.; et. al.; *Gazzetta Chimica Italiana*, 116, 649-652, (1986); (3) King, J. F. and Hillhouse, J. H.; *Can. J. Chem.*; 54, 498, (1976); and (4) Szymonifka, M. J. and Heck, J. V.; *Tet. Lett.*; 30, 2869-2872, (1989).

[0120] A detailed description of the conversion of a compound of Formula IIa to a compound of Formula (I) where n is 1 and E is -N(R⁶)SO₂- is given in Example 7.

[0121] Compounds of Formula (I) where E is -N(R⁶)SO₂N(R⁵)- are prepared as described in Scheme I below:


Scheme I



[0122] A compound of Formula (I) where E is a sulfamide group can be prepared by reacting a compound of Formula II (a-c) with a sulfamoyl halide, utilizing the reaction conditions described in method (i) of Scheme E. Sulfamoyl halides are commercially available or may be prepared by methods such as those described in Graf, R; *German Patent*, 931225 (1952) and Catt, J.D. and Matler, W.L.; *J. Org. Chem.*, 39, 566-568, (1974).

[0123] Compounds of Formula (I) where E is -N(R⁶)C(O)- are prepared as described in Scheme J below:

Scheme J



[0124] A compound of Formula (I) where E is an inverse amide can be prepared by reacting a compound of Formula II(d-f) with an amine in the presence of a suitable coupling agent (e.g., N,N-dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide, and the like) in a suitable organic solvent such as methylene chloride, tetrahydrofuran, dimethylformamide and the like. A detailed description of a compound of Formula II(d) to a compound of Formula I where n is 1 and E is -NHC(O)- is given in Example 6.

[0125] The compounds of the invention are CCR-3 receptor antagonists and inhibit eosinophil recruitment by CCR-3 chemokines such as RANTES, eotaxin, MCP-2, MCP-3 and MCP-4. Compounds of this invention and compositions containing them are useful in the treatment of eosinophil-induced diseases such as inflammatory or allergic diseases and including respiratory allergic diseases such as asthma, allergic rhinitis, hypersensitivity lung diseases, hypersensitivity pneumonitis, eosinophilic pneumonias (e.g., chronic eosinophilic pneumonia); inflammatory bowel diseases (e.g., Crohn's disease and ulcerative colitis); and psoriasis and inflammatory dermatoses such as dermatitis and eczema.

[0126] The CCR-3 antagonistic activity of the compounds of this invention was measured by *in vitro* assays such as ligand binding and chemotaxis assays as described in more detail in Examples 15, 16 and 17. *In vivo* activity was assayed in the Ovalbumin induced Asthma in Balb/c Mice Model as described in more detail in Example 18.

[0127] In general, the compounds of this invention will be administered in a therapeutically effective amount by any of the accepted modes of administration for agents that serve similar utilities. The actual amount of the compound of this invention, i.e., the active ingredient, will depend upon numerous factors such as the severity of the disease to be treated, the age and relative health of the subject, the potency of the compound used, the route and form of administration, and other factors.

[0128] Therapeutically effective amounts of compounds of Formula (I) may range from approximately 0.05-20 mg per kilogram body weight of the recipient per day; preferably about 0.1-10 mg/kg/day. Thus, for administration to a 70 kg person, the dosage range would most preferably be about 7 mg to 0.7 g per day.

[0129] In general, compounds of this invention will be administered as pharmaceutical compositions by any one of the following routes: oral, inhalation (e.g., intranasal or oral inhalation) or parenteral (e.g., intramuscular, intravenous or subcutaneous) administration. A preferred manner of administration is oral using a convenient daily dosage regimen which can be adjusted according to the degree of affliction. Compositions can take the form of tablets, pills, capsules, semisolids, powders, sustained release formulations, solutions, suspensions, elixirs, or any other appropriate compositions. Another preferred manner for administering compounds of this invention is inhalation. This is an effective means for delivering a therapeutic agent directly to the respiratory tract for the treatment of diseases such as asthma and other similar or related respiratory tract disorders (see U. S. Patent 5,607,915).

[0130] The choice of formulation depends on various factors such as the mode of drug administration and the bioavailability of the drug substance. For delivery via inhalation the compound can be formulated as liquid solutions or suspensions, aerosol propellants or dry powder and loaded into a suitable dispenser for administration. There are three types of pharmaceutical inhalation devices- nebulizer inhalers, metered-dose inhalers (MDI) and dry powder inhalers (DPI). Nebulizer devices produce a stream of high velocity air that causes the therapeutic agents (which has been formulated in a liquid form) to spray as a mist which is carried into the patient's respiratory tract. MDI's typically have the formulation packaged with a compressed gas. Upon actuation, the device discharges a measured amount of therapeutic agent by compressed gas, thus affording a reliable method of administering a set amount of agent. DPI's administer therapeutic agents in the form of a free flowing powder that can be dispersed in the patient's inspiratory air-stream during breathing by the device. In order to achieve a free flowing powder, the therapeutic agent is formulated with an excipient, such as lactose. A measured amount of the therapeutic is stored in a capsule form and is dispensed to the patient with each actuation. Recently, pharmaceutical formulations have been developed especially for drugs that show poor bioavailability based upon the principle that bioavailability can be increased by increasing the surface area i.e., decreasing particle size. For example, U.S. Pat. No. 4,107,288 describes a pharmaceutical formulation having particles in the size range from 10 to 1,000 nm in which the active material is supported on a crosslinked matrix of macromolecules. U.S. Pat. No. 5,145,684 describes the production of a pharmaceutical formulation in which the drug substance is pulverized to nanoparticles (average particle size of 400 nm) in the presence of a surface modifier and then dispersed in a liquid medium to give a pharmaceutical formulation that exhibits remarkably high bioavailability.

[0131] The compositions are comprised of in general, a compound of Formula (I) in combination with at least one pharmaceutically acceptable excipient. Acceptable excipients are non-toxic, aid administration, and do not adversely affect the therapeutic benefit of the compound of Formula (I). Such excipient may be any solid, liquid, semi-solid or, in the case of an aerosol composition, gaseous excipient that is generally available to one of skill in the art.

[0132] Solid pharmaceutical excipients include starch, cellulose, talc, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, magnesium stearate, sodium stearate, glycerol monostearate, sodium chloride, dried skim milk and the like. Liquid and semisolid excipients may be selected from glycerol, propylene glycol, water, ethanol and various oils, including those of petroleum, animal, vegetable or synthetic origin, e.g., peanut oil, soybean oil, mineral oil, sesame oil, etc. Preferred liquid carriers, particularly for injectable solutions, include water, saline, aqueous dextrose, and glycols.

[0133] Compressed gases may be used to disperse a compound of this invention in aerosol form. Inert gases suitable for this purpose are nitrogen, carbon dioxide, etc.

[0134] Other suitable pharmaceutical excipients and their formulations are described in *Remington's Pharmaceutical Sciences*, edited by E. W. Martin (Mack Publishing Company, 18th ed., 1990).


[0135] The level of the compound in a formulation can vary within the full range employed by those skilled in the art. Typically, the formulation will contain, on a weight percent (wt%) basis, from about 0.01-99.99 wt% of a compound of Formula (I) based on the total formulation, with the balance being one or more suitable pharmaceutical excipients. Preferably, the compound is present at a level of about 1-80 wt%. Representative pharmaceutical formulations containing a compound of Formula (I) are described in Example 14.

ExamplesSynthetic Examples

5 Example 1

Synthesis of N-{1(*S*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide dihydrochloride salt

10 [0136]

25 Step 1

[0137] 3,4-Dichlorobenzyl bromide (35.20 g, 150 mmol) was added to a solution of N-(*tert*-butoxycarbonyl)piperazine (24.84 g, 130 mmol) and triethylamine (20.91 ml, 150 mmol) in chloroform (100 ml) over 30 min. After 1 h, the reaction mixture was diluted with ethyl acetate and the product was precipitated out as the hydrochloride salt by adding 1N aqueous hydrogen chloride solution. The solid product was filtered, washed with water and then resuspended in ethyl acetate. Two equivalents of 1N aqueous sodium hydroxide solution was added and the free amine was extracted into ethyl acetate. The organic layer was separated, dried over magnesium sulfate, filtered and concentrated to provide 1-(*tert*-butoxycarbonyl)-4-(3,4-dichlorobenzyl)piperazine (45 g).

35 Step 2

[0138] Trifluoroacetic acid (75 ml, 974 mmol) was added to a solution 1-(*tert*-butoxycarbonyl)-4-(3,4-dichlorobenzyl)piperazine (45 g, 130 mmol) in chloroform (75 ml). The reaction mixture was stirred for 1 h at room temperature and then made basic with a sodium hydroxide solution. The product was extracted into ethyl acetate and the organic layer was washed with sodium bicarbonate solution, dried over magnesium sulfate and concentrated *in vacuo* to give 1-(3,4-dichlorobenzyl)piperazine (35.8 g) as a solid.

45 Step 3

[0139] 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (5.08 g, 26.5 mmol) was added to a solution of 1-(3,4-dichlorobenzyl)piperazine (5 g, 20.4 mmol) and L-Boc-valine (5.76g, 26.5 mmol) in methylene chloride. After 2 h, the product was extracted into ethyl acetate. The organic layer was washed with sodium bicarbonate solution, dried over magnesium sulfate, filtered and concentrated *in vacuo*. Column chromatography with hexanes/ethyl acetate (1:1) as the eluant gave 1-(*S*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylcarbonyl]-N-(*tert*-butoxycarbonyl)-2-methylpropylamine (5.46 g) as a foam.

55 Step 4

[0140] 1 N ethereal hydrogen chloride solution (80 ml, 80 mmol) was added to a solution of 1(*S*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylcarbonyl]-N-(*tert*-butoxycarbonyl)-2-methylpropylamine (4.28 g, 9.64 mmol) in methanol (50 ml) and the reaction mixture was heated at 70 °C. After 2.5 h, the reaction mixture was concentrated and the solid was suspended in ether and filtered to give 1-(*S*)-[4-(3,4-dichlorobenzyl)piperazin-1-yl carbonyl]-2-methylpropylamine as the

bis hydrochloride salt. The product was dissolved in water, treated with triethylamine (4.0 ml, 28.9 mmol) and the free amine was extracted into ethyl acetate. The ethyl acetate layer was dried over magnesium sulfate, filtered and concentrated to give 1-(S)-[4-(3,4-dichlorobenzyl)piperazin-1-ylcarbonyl]-2-methylpropylamine (3.2 g) as the free amine.

5 Step 5

[0141] A 1.0 M diborane solution in tetrahydrofuran (65.2 ml, 65.2 mmol) was added to a solution of 1-(S)-[4-(3,4-dichlorobenzyl)piperazin-1-ylcarbonyl]-2-methylpropylamine (3.2 g, 9.3 mmol) in tetrahydrofuran (15 ml). The mixture was heated at reflux under nitrogen atmosphere for 2 h and then concentrated *in vacuo*. The residue was dissolved in methanol, acidified with 6 N hydrogen chloride solution (50 ml) and then reheated to 70 °C. After heating for 1 h, the reaction mixture was cooled and basified with a sodium hydroxide solution and the product was extracted into ethyl acetate. The ethyl acetate layer was washed with sodium bicarbonate solution, dried over magnesium sulfate, filtered and concentrated to provide 1-(S)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropylamine (3.53 g) as an oil.

15 Step 6

[0142] *p*-Toluoyl chloride (0.48 ml, 3.63 mmol) was added to a solution of 1(S)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropylamine (1 g, 3.0 mmol) and triethylamine (0.633 ml, 4.54 mmol) in methylene chloride under nitrogen atmosphere. After 1h, the product was extracted into ethyl acetate and the organic layer was washed with sodium bicarbonate solution, dried over magnesium sulfate, filtered and concentrated *in vacuo*. Column chromatography with hexanes/ ethyl acetate (1:2) gave N-{1(S)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide (1.2 g) as an oil. The free amine was dissolved in ether and 3.5 equivalents of 1N ethereal HCl solution (10.7 ml) was added. Filtration of the resulting solid provided N-{1(S)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide (1.2 g) as the bis hydrochloride salt, mp 227.8-228.9 °C.

1. Proceeding as described in Example 1, Steps 1-4 above but substituting L-Boc-valine in Step 3 with DL-Boc-valine gave 1-(RS)-[4-(3,4-dichlorobenzyl)piperazin-1-ylcarbonyl]-2-methylpropylamine which upon reduction (step 5) and reaction with 3,4-methylenedioxybenzoyl chloride (step 6) gave N-{1(RS)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-3,4-methylenedioxybenzamide.
2. Proceeding as described in Example 1, Steps 1-4 above but substituting L-Boc-valine in Step 3 with L-Boc-*tert*-leucine (commercially available from Fluka) gave 1-(S)-[4-(3,4-dichlorobenzyl)piperazin-1-ylcarbonyl]-2,2-dimethylpropylamine which was reduced to 1-(S)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2,2-methylpropylamine (Step 5) and then reacted with:

- 35 4-Methylsulfonylbenzoyl chloride;
4-Acetoxybenzoyl chloride;
4-N,N-Dimethylaminobenzoyl chloride;
5-Methyl-2-thenoyl chloride; and
4-Methylbenzoyl chloride, to give
- 40 N-{1(S)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-4-methylsulfonylbenzamide dihydrochloride salt, mp 190-191 °C;
N-{1(S)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-4-acetoxybenzamide dihydrochloride salt, mp 241-242 °C;
N-{1(S)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-4-N,N-dimethylaminobenzamide dihydrochloride salt, mp 101.5-105 °C;
- 45 N-{1(S)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-5-methyl-2-thiophenecarboxamide dihydrochloride salt, mp 249-253 °C; and
N-{1(S)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-4-methylbenzamide dihydrochloride salt by following the procedure described in Steps 5 and 6 above.

- 50 3. Proceeding as described in Example 1, Steps 1-4 above but substituting L-Boc-valine in Step 3 with DL-Boc-*tert*-leucine gave 1-(RS)-[4-(3,4-dichlorobenzyl)piperazin-1-ylcarbonyl]-2,2-methylpropylamine which was reduced to 1-(RS)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2,2-methylpropylamine (Step 5) and reacted with:

- 55 3-Cyanobenzoyl chloride; and
3,4-Difluorobenzoyl chloride, to give
N-{1(RS)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-3-cyanobenzamide; and
N-{1(RS)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-3,4-difluorobenzamide, respec-


tively.

4. Proceeding as described in Example 1, but substituting N-(*tert*-butoxycarbonyl)-piperazine in Step 1 with 3-methyl-N-(*tert*-butoxycarbonyl)piperazine and L-Boc-valine in Step 3 with DL-Boc-valine gave N-{1(*RS*)-[3-methyl-4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide dihydrochloride salt.

Example 2

Synthesis of 1-{2-[4-(3,4-Dichlorobenzyl)-piperazin-1-yl]cyclohexyl}-3-(3-methoxyphenyl)urea

[0143]



Step 1

[0144] A mixture of 1-nitrocyclohexene (311 mg, 2.45 mmol), dichlorobenzylpiperazine [prepared as described in Example 1] (600 mg, 2.45 mmol), and triethylamine (512 μ l, 3.68 mmol) in methylene chloride (10 ml) was stirred at room temperature under nitrogen atmosphere for 17 h. The reaction mixture was diluted with ethyl acetate, washed with an aqueous sodium bicarbonate solution, dried over magnesium sulfate and concentrated *in vacuo*. The residue was chromatographed with hexanes/ ethyl acetate (2: 1) as the eluant to provide 1-(3,4-dichlorobenzyl)-4-(2-nitrocyclohexyl)piperazine (461 mg) as an oil.

Step 2


[0145] Platinum (IV) oxide (15 mg, 0.07 mmol) was added to a solution of 1-(3,4-dichlorobenzyl)-4-(2-nitrocyclohexyl)piperazine (187 mg, 0.50 mmol) in glacial acetic acid (5 ml). The reaction mixture was stirred under hydrogen atmosphere at room temperature for 16 h. The reaction mixture was then filtered through a pad of celite. The filtrate was made basic with a 15% sodium hydroxide solution and the product was extracted into ethyl acetate. The organic layer was dried over magnesium sulfate, filtered and concentrated *in vacuo* to provide 1-(3,4-dichlorobenzyl)-4-(2-aminocyclohexyl)piperazine (119 mg) as an oil.

[0146] A solution of 1-(3,4-dichlorobenzyl)-4-(2-aminocyclohexyl)piperazine (100 mg, 0.29 mmol) and 3-methoxyphenyl isocyanate (38 μ l, 0.29 mmol) in methylene chloride (5 ml) was stirred at room temperature. After 17 h, the product was extracted into ethyl acetate, washed with a bicarbonate solution, dried over magnesium sulfate and concentrated *in vacuo*. Column chromatography with methylene chloride/ methanol (97: 3) as the eluant gave 1-{2-[4-(3,4-dichlorobenzyl)piperazin-1-yl]cyclohexyl}-3-(3-methoxyphenyl)urea (64 mg) as an oil consisting of the trans diastereomer.

Example 3

Synthesis of N-{1-(S)-[4-(3,4-Dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide

[0147]



Step 1

[0148] *n*-Butyllithium (43.2 ml, 2M in pentane, 108 mmol) was slowly added to an ice-cooled suspension of 3,4-dichlorobenzyl triphenylphosphonium bromide (54 g, 108 mmol) (prepared by stirring equimolar amounts of 3,4 dichlorobenzyl bromide and triphenylphosphine in THF at 65° overnight) in dry THF (500 ml) under an argon atmosphere. After 15 min., the reaction mixture was allowed to warm to room temperature, and was stirred for an additional 2 h. 1-*tert*-butoxycarbonyl-4-piperidone (21.42 g, 108 mmol) was added and the stirring was continued overnight. Hexane (2 l) was added and the reaction was stirred and then filtered. The filtrate was concentrated *in vacuo* to give 41.8 g of an orange gum. Column chromatography on 0.5 kg flash grade silica, eluting with a gradient of 70% methylene chloride/hexane through 100 % methylene chloride, followed by a gradient of 1% methanol/methylene chloride through 5% methanol /methylene chloride gave 1-(*tert*-butoxycarbonyl)-4-(3,4-dichloro-benzylidene)piperidine (29 g) as a light tan oil.

Step 2

[0149] Platinum oxide (0.3 g) was added to a solution of 1-(*tert*-butoxycarbonyl)-4-(3,4-dichlorobenzylidene)piperidine (29 g, 84.7 mmol) in ethyl acetate (500 ml) and the mixture was stirred under a hydrogen atmosphere overnight. The reaction mixture was filtered through celite and the filtrate was concentrated to give 1-(*tert*-butoxycarbonyl)-4-(3,4-dichlorobenzyl)piperidine (30 g) as a tan oil.

Step 3

[0150] Trifluoroacetic acid (50 ml) was added to a solution of 1-(*tert*-butoxycarbonyl)-4-(3,4-dichlorobenzyl)piperidine (24 g, 69.7 mmol) in methylene chloride (150 ml) and the reaction mixture was stirred for 1 h. The solvent was removed under reduced pressure, followed by addition of ethyl acetate (200 ml), and the resulting mixture was made basic with 1N aqueous sodium hydroxide. The organic layer was separated, dried over magnesium sulfate and the solvent was removed under reduced pressure to give 4-(3,4-dichlorobenzyl)piperidine (17.1 g) as light brown solid.

Step 4

[0151] L-BOC-Valine (1.3 g, 5.98 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (1.15 g, 5.98 mmol) were added to a solution of 4-(3,4-dichlorobenzyl)piperidine (1.12 g, 4.57 mmol) in methylene chloride (15 ml) and the reaction mixture was stirred at room temperature under an argon atmosphere. After 3 h the solvents were removed under *vacuo* and water (10 ml) and ethyl acetate (25 ml) were added. The organic layer was separated, dried over magnesium sulfate and concentrated under reduced pressure. Column chromatography with 15-20% ethyl acetate /hexane as the eluant gave 1-(S)-[4-(3,4-dichlorobenzyl)-piperidin-1-ylcarbonyl]-N-(*tert*-butoxycarbonyl)-2-methylpropylamine (1.89 g) as a gummy foam.

Step 5

[0152] To a solution of 1-(*S*)-[4-(3,4-dichlorobenzyl)piperidin-1-ylcarbonyl]-*N*-(*tert*-butoxycarbonyl)-2-methylpropylamine (5.9g, 13.2 mmol) in methylene chloride (100 ml) was added trifluoroacetic acid (30 ml) at room temperature. After 4 h, the reaction mixture was concentrated and the residue was stirred with ethyl acetate (200 ml) and water (100 ml) while adjusting the pH to 8 with 15% aqueous sodium hydroxide solution. The organic layer was separated and the aqueous layer was extracted twice with ethyl acetate. The combined organic portions were dried over magnesium sulfate, filtered and concentrated *in vacuo* to give 1-(*S*)-[4-(3,4-dichlorobenzyl)piperidin-1-ylcarbonyl]-2-methylpropylamine (4.53g) as a colorless gum.

Step 6


[0153] To a solution of 1-(*S*)-[4-(3,4-dichlorobenzyl)piperidin-1-ylcarbonyl]-2-methylpropylamine (4.53 g, 13.3 mmol) in dry tetrahydrofuran (100 ml) was added diborane (92.4 ml, 92.4 mmol, 1 M in THF) and the reaction mixture was stirred under argon at 65 °C. After 3 h, the reaction mixture was cooled in an ice bath and aqueous hydrochloric acid (60 ml, 6 N) was slowly added with stirring. The reaction mixture was concentrated on a rotovap and the aqueous solution was stirred at 100 °C. After 1 h, the reaction mixture was cooled to 0 °C and potassium hydroxide pellets were slowly added until pH 8 was obtained. The solution was extracted twice with ethyl acetate (100 ml), dried over magnesium sulfate and concentrated *in vacuo*. The colorless liquid (3.84 g) was flash chromatographed, eluting with 2.5-10% MeOH/CH₂Cl₂ containing 1% NH₄OH. The free amine was dissolved in anhydrous ether and ethereal HCl was added to afford 1-(*S*)-[4-(3,4-dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropylamine as the HCl salt.

Step 7

[0154] *p*-Toluoyl chloride (0.14 ml, 1 mmol) was added to a solution of 1-(*S*)-[4-(3,4-dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropylamine (0.33 g, 1 mmol) in dry pyridine (7 ml) at 0 °C under argon atmosphere. The solution was stirred for 30 min., at 0 °C and then allowed to warm to room temperature. After 3 h, the reaction mixture was concentrated under reduced pressure and water (10 ml) was added. The product was extracted with ethyl acetate and the organic layer was dried with magnesium sulfate and then concentrated *in vacuo*. The crude product (0.5 g) was chromatographed by eluting with a gradient of from 1% to 4% methanol / methylene chloride to give *N*-[1-(*S*)-[4-(3,4-dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl]-4-methylbenzamide (0.25 g) as partially solidified gum.

Example 4

N-[1-(*RS*)-[4-(4-Amino-5-chloro-2-methoxyphenylcarbonylaminomethyl)piperidin-1-yl-methyl]-2-methylpropyl]-4-methylbenzamide

[0155]Step 1

[0156] Diisopropylethyl amine (17.4 ml, 134 mmol) was added a solution of (*DL*)-valinol (9.85 g, 95 mmol) in methylene chloride (100 ml). The reaction mixture was cooled to 0 °C, treated with a solution of *p*-toluoyl chloride (12.8 ml, 91 mmol) in methylene chloride (50 ml) and then allowed to warm to room temperature. After stirring for 3 hours, excess aqueous sodium hydroxide solution was added and the reaction was transferred to a separatory funnel. The organic layer was separated and the aqueous layer washed with one portion of methylene chloride. The combined organic layer

ers were washed with water and brine, dried over magnesium sulfate and concentrated *in vacuo*. Chromatography eluting with 25% ethyl acetate in hexanes, followed by 50% ethyl acetate in hexanes gave N-*p*-toluoyl valinol (18.04 g).

Step 2

[0157] Dimethylsulfoxide (2.2 ml, 31 mmol) was slowly added via syringe to a stirred -78 °C solution of oxalyl chloride (15 ml, 171 mmol) in methylene chloride (35 ml) under inert atmosphere. After 10 min., a solution of N-*p*-toluoyl valinol (6.0 g, 29 mmol) in methylene chloride (50 ml) was added and the stirring was continued for additional 15 min. Triethylamine was added (6 ml, 389 mmol) and the reaction was allowed to warm to ambient temperature. After 1.5 h, the reaction was diluted with 50% ethyl acetate in hexanes and washed with water and brine. Filtration through a pad of silica gel and subsequent solvent removal left a solid residue. Chromatography eluting with 20% ethyl acetate in hexanes, then 33% ethyl acetate in hexanes gave N-*p*-toluoyl valinaldehyde (3.6 g) as a solid, which was utilized in Step 6.

Step 3

[0158] *tert*-Butyloxycarbonyl anhydride (6.69g, 30.6 mmol) was added to a solution of 4-(aminomethyl)piperidine (7 g, 61.3 mmol) in chloroform (40 ml) at 0 °C. The reaction mixture was allowed to warm to room temperature over 3 h and then stirred an additional 15 h. The reaction mixture was washed with water, the organic layer was separated and dried over magnesium sulfate, filtered and concentrated to provide N-*tert*-butyloxy-carbonyl-4-(aminomethyl)piperidine (6.55 g) as a pale yellow oil.

Step 4

[0159] 1,1'-Carbonyldiimidazole (1.61 g, 9.91 mmol) was added to a solution of 4-amino-5-chloro-2-methoxybenzoic acid (2.0 g, 9.91 mmol) in dimethylformamide (5 ml). After stirring for 5 min. at room temperature, solution of N-*tert*-butyloxycarbonyl-4-(aminomethyl)piperidine (1.77 g, 8.26 mmol) in dimethylformamide (5 ml) was added and the reaction mixture was heated at 55 °C. After 23 h, the product was extracted into ethyl acetate and the organic layer was washed with aqueous sodium bicarbonate, dried over magnesium sulfate, filtered and concentrated. Flash chromatography of the crude product with 2: 1, ethyl acetate/hexanes as the eluant gave N-*tert*-butyloxycarbonyl-4-(4-amino-5-chloro-2-methoxyphenylcarbonylaminomethyl)piperidine (2.63 g) as a foam.

Step 5

[0160] To a solution of N-*tert*-butyloxycarbonyl-4-(4-amino-5-chloro-2-methoxyphenyl-carbonylaminomethyl)piperidine (500 mg, 1.26 mmol) in methanol solution (30 ml) was added an ethereal 1N hydrogen chloride solution (12.6 ml, 12.6 mmol). The solution was heated at 50 °C. The reaction mixture became heterogeneous within 5 min. After 1.5 h, the reaction mixture was concentrated to give a white solid which was suspended in diethylether and filtered to give 4-(4-amino-5-chloro-2-methoxyphenylcarbonylamino-methyl)piperidine as a bis-hydrochloride salt (433 mg).


Step 6

[0161] To a heterogeneous mixture of 4-(4-amino-5-chloro-2-methoxyphenylcarbonyl-aminomethyl)piperidine (304 mg, 0.82 mmol) and triethylamine (309 µl, 2.21 mmol) in methylene chloride (20 ml) was added aldehyde N-*p*-toluoyl valinaldehyde (150 mg, 0.68 mmol) and sodium triacetoxymethylborohydride (216 mg, 1.02 mmol) at room temperature. After 1 h the reaction mixture was diluted with ethyl acetate and washed with sodium bicarbonate solution. The organic layer was separated, dried over magnesium sulfate and concentrated *in vacuo*. Flash chromatography of the crude product with 97:3, methylene chloride/ methanol as the eluant gave N-{1(*RS*)-[4-(4-amino-5-chloro-2-methoxyphenylcarbonylaminomethyl)piperidin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide (139 mg) as a foam.

Example 5

{1-(4-Methylbenzoyl)-2(*R*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]}-pyrrolidine dihydrochloride salt

[0162]



Step 1

[0163] 1-Ethyl-3-(3'-dimethylaminopropyl)carbodiimide hydrochloride (1.03 g, 7 mmol) was added to a solution of N-BOC-*D*-Proline (1.51 g, 7 mmol) in methylene chloride (12 ml) and the reaction mixture was stirred at room temperature. After 0.5 h, 3,4-dichlorobenzylpiperazine (1.32 g, 5.4 mmol) [prepared as described in Example 1] was added and the stirring was continued for 16 h. The reaction mixture was then quenched with water, basified with saturated sodium bicarbonate solution and extracted with ethyl acetate. The ethyl acetate layer was separated and washed with brine, dried over magnesium sulfate and concentrated. Flash chromatography with 30% acetone in dichloromethane as the eluant gave 2(*R*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylcarbonyl]-N-(*tert*-butoxycarbonyl) pyrrolidine (1.2 g) as an oil.

Step 2

[0164] Borane (15.8 ml, 1.0 M solution in THF) was added dropwise to a solution of 2(*R*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylcarbonyl]-N-(*tert*-butoxycarbonyl) pyrrolidine (1.0 g, 2.26 mmol) in tetrahydrofuran (5 ml) was under nitrogen atmosphere and the reaction mixture was heated at reflux. After 2 h, the reaction mixture was cooled to room temperature and quenched with 6 N hydrochloric acid. The reaction mixture was re-heated at reflux for another 2 h, cooled to room temperature and then basified with 10% aqueous sodium hydroxide solution. The product was extracted into ethyl acetate and the organic layer was separated and washed with brine, dried over magnesium sulfate, and concentrated. Purification with column chromatography with methylene chloride containing 10% ammonium hydroxide in methanol as the eluant gave 2(*R*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]pyrrolidine (0.62 g) as an oil.


Step 3

[0165] Triethylamine (0.19 ml, 1.5 mmol) and *p*-toluoyl chloride (0.11 ml, 0.84 mmol) were added to a solution of 2(*R*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]pyrrolidine (0.25 g, 0.76 mmol) in methylene chloride (5 ml). After 1 h, the reaction mixture was quenched with water and the product was extracted into ethyl acetate. The ethyl acetate layer was washed with brine, dried over magnesium sulfate and concentrated. Preparative TLC with 5% methanol in methylene chloride afforded {1-(4-methylbenzoyl)-2(*R*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]}pyrrolidine 0.245 g of an oil which was converted to its dihydrochloride salt and recrystallized from a methanol-ethyl acetate mixture to give {1-(4-methylbenzoyl)-2(*R*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]}pyrrolidine dihydrochloride (0.18 g) as a white solid; m.p. 249.6-250.1 °C.

Example 6

Synthesis of 2-(*RS*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-N-(4-methylphenyl)propionamide

[0166]

Step 1

[0167] A mixture of 3,4-dichlorobenzylpiperazine (2.47 g, 10.1 mmol), methyl methacrylate (2.2 ml, 21 mmol), and sodium methoxide (49 mg, 0.91 mmol) in methanol (20 ml) was heated to reflux. After 72 h, the reaction mixture was concentrated under reduced pressure to yield pale-yellow oil (3.02 g). Column chromatography on silica gel with ethyl acetate as eluant afforded methyl 2-(*RS*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]propionate (1.52 g) as a pale-yellow oil.

Step 2

[0168] Methyl 2-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]propionate (470 mg, 1.36 mmol) and lithium hydroxide monohydrate (185 mg, 4.41 mmol) were dissolved in water (5 ml) and methanol (15 ml) and stirred at room temperature. After 21 h, the solvents were removed under reduced pressure. The resulting residue was taken up in a mixture of methylene chloride and water, and the aqueous layer was made acidic (~ pH 4) with 1M HCl. The layers were separated and the aqueous phase was extracted with several portions of methylene chloride. The combined organic layers were dried over sodium sulfate and concentrated under reduced pressure to afford 2-(*RS*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]propionic acid (444 mg) as a pale-yellow foam.


Step 3

[0169] 1-Ethyl-3-(3'-dimethylaminopropyl)carbodiimide hydrochloride (388 mg, 2.02 mmol) was added to a solution of 2-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-propionic acid (609 mg, 1.84 mmol) *p*-toluidine (219 mg, 2.04 mmol) and dimethylaminopyridine (22 mg, 0.18 mmol) in methylene chloride (20 ml). After 4 h, the mixture was diluted with methylene chloride and washed with water. The organic layer was separated, dried with sodium sulfate and the solvent removed under reduced pressure to give an orange foam (839 mg). Column chromatography on silica gel with ethyl acetate as eluant yielded 2-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-N-(4-methyl-phenyl)propionamide (636 g, 82%) as a white foam. A 1M solution of hydrochloric acid in ether (4.2 ml) was added to a solution of this product (593 mg, 1.41 mmol) in methanol and the solvent was removed under reduced pressure. The resulting viscous oil was triturated with ether/hexanes to yield 2-(*RS*)-[4-(3,4-dichlorobenzyl)piperazin-1-yl-methyl]-N-(4-methylphenyl)propionamide (245 mg) as the hydrochloride salt, m.p. 256.2-256.7 °C.

Example 7

N-{1-(*R*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-fluorobenzenesulfonamide dihydrochloride salt

[0170]




[0171] Triethylamine (21 ml, 0.15 mmol) and 4-fluorobenzenesulfonyl chloride (29 mg, 0.15 mmol) were added to a solution of 1-(*R*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropylamine (50 mg, 0.15 mmol) [prepared as described in Example 1 above] in methylene chloride (1.5 ml). The reaction mixture was stirred for 2 h, then concentrated. Purification by flash chromatography with 20:1 methylene: methanol as the eluant gave N-{1-(*R*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-fluorobenzenesulfonamide (49 mg) as a solid.

Example 8

1-{1-(*RS*)-[4-(3,4-Dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-3-methoxybenzyl-2-thiourea

[0172]




[0173] 3-Methoxybenzyl isothiocyanate (22 mg, 0.12 mmol) was added to a solution of 1-(*RS*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropylamine (40 mg, 0.12 mmol) [prepared as described in Example 1 by substituting *L*-N-BOC-valine with *DL*-N-BOC-valine] in methylene chloride (1.5 ml). The reaction mixture was stirred for 2 h and then concentrated. Purification by flash chromatography with 20:1 methylene chloride: methanol as the eluant gave 1-{1-(*RS*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-3-methoxybenzyl-2-thiourea (18 mg) as a solid.

Example 9

1-{1(*RS*)-[1-(3,4-Dichlorobenzyl)piperidin-4-yl]-2-methylpropyl}-3-(3-methoxyphenyl)urea

[0174]



Step 1

[0175] 3,4-Dichlorobenzyl bromide (1.83 g 7.63 mmol) was added to a solution of ethyl isonipecotate (1 g, 6.36 mmol) and triethylamine (1.33 ml, 9.54 mmol) in methylene chloride solution (10 ml) at room temperature. After 17 h, the reaction mixture was diluted with ethyl acetate and washed with aqueous sodium bicarbonate solution (10 ml). The organic layer was dried over magnesium sulfate, filtered, and concentrated. Flash chromatography with 9: 1, hexanes/ ethyl acetate followed by 6: 1 hexanes/ ethyl acetate gave ethyl 1-(3,4-dichlorobenzyl)isonipecotate (1.78 g) as a pale yellow oil.

Step 2

[0176] A 1.5 M solution of DIBAL-H™ in toluene (1.27 ml, 1.90 mmol) was added dropwise over 10 min. to a solution of ethyl 1-(3,4-dichlorobenzyl)isonipecotate (500 mg, 1.58 mmol) in toluene solution (5 ml) at -78 °C. After 40 min., aqueous sodium bicarbonate solution was added and the reaction mixture was warmed to room temperature. The product was extracted into diethyl ether. The organic layer was dried over magnesium sulfate, filtered and concentrated to provide 1-(3,4-dichlorobenzyl)-4-formylpyridine (430 mg) as an oil.

Step 3

[0177] Isopropylmagnesium chloride in tetrahydrofuran (3.7 ml, 7.35 mmol, 2.0 M) was added to a solution of 1-(3,4-dichlorobenzyl)-4-formylpyridine (1 g, 3.67 mmol) in tetrahydrofuran (10 ml) at 0 °C. After 30 min., the reaction mixture was warmed to room temperature and stirred an additional 30 min. The reaction mixture was then diluted with diethyl ether and washed with sodium bicarbonate solution (10 ml). The organic layer was separated and dried over magnesium sulfate, filtered and concentrated. The residue was subjected to flash chromatography (1: 1, hexanes/ ethyl acetate) to provide alcohol 1-(*RS*)-[(3,4-dichlorobenzyl)pyridin-4-yl]-2-methylpropanol (617 mg) as an oil.

Step 4

[0178] Pyridinium dichromate (6 g, 15.9 mmol) was added to a solution of 1-(*RS*)-[(3,4-dichlorobenzyl)pyridin-4-yl]-2-methylpropanol (617 mg, 1.95 mmol) in N,N-dimethylformamide (35 ml) at 0 °C. After 4.5 h, the reaction mixture was diluted with ethyl acetate and washed with sodium bicarbonate solution (30 ml). The organic layer was separated and dried over magnesium sulfate, filtered and concentrated. The residue was subjected to flash chromatography (1.5: 1, hexanes/ ethyl acetate) to provide 1-[(3,4-dichlorobenzyl)pyridin-4-yl]-2-methylpropanone (432 mg) as an oil.

Step 5

[0179] Ammonium acetate (1.43 g, 18.33 mmol) followed by sodium cyanoborohydride (130 mg, 1.86 mmol) were added to a solution of 1-[(3,4-dichlorobenzyl)pyridin-4-yl]-2-methylpropanone (583 mg, 1.86 mmol) in methanol (10 ml) at room temperature. After 50 h, additional amounts of ammonium acetate (1.43 g, 18.33 mmol) and sodium cyanoborohydride (130 mg, 1.86 mmol) were added. After 17 h, the reaction mixture was diluted with ethyl acetate and

washed with sodium bicarbonate solution (10 ml). The organic layer was separated and dried over magnesium sulfate, filtered and concentrated. The residue was dissolved in ethyl acetate and the product extracted into 1N HCl solution. The acidic aqueous layer was made basic with sodium hydroxide solution and the product was extracted into ethyl acetate. The organic layer was dried over magnesium sulfate, filtered and concentrated. The residue was subjected to flash chromatography (90: 10, methylene chloride/ methanol followed by 80: 20) to provide 1(*RS*)-[(3,4-dichlorobenzyl)pyridin-4-yl]-2-methylpropylamine (390 mg) as an oil.

Step 6


[0180] 3-methoxyphenyl isocyanate (37 μ l, 0.28 mmol) was added to a solution of provide 1(*RS*)-[(3,4-dichlorobenzyl)pyridin-4-yl]-2-methylpropylamine (88 mg, 0.28 mmol) in methylene chloride solution (2 ml) at room temperature. After 1 h, the reaction mixture was diluted with ethyl acetate and washed with sodium bicarbonate solution (5 ml). The organic layer was separated and dried over magnesium sulfate, filtered and concentrated to provide 1-{1(*RS*)-[1-(3,4-dichlorobenzyl)piperidin-4-yl]-2-methylpropyl}-3-(3-methoxyphenyl)urea. (104 mg) as a white solid. mp 161.4-161.8 °C.

[0181] Proceeding as described in Example 9, Step 6 above but substituting 1(*RS*)-[(3,4-dichlorobenzyl)pyridin-4-yl]-2-methylpropylamine with 1-(*R*)-[4-(3,4-dichlorobenzyl)-piperazin-1-ylmethyl]-2-methylpropylamine (prepared as described in Example 1) gave 1-{1(*R*)-[1-(3,4-dichlorobenzyl)piperazin-1-yl]-2-methylpropyl}-3-(3-methoxyphenyl)urea.

Example 10

N-{1-[1(*RS*)-[1-(3,4-Dichlorobenzyl)piperidin-4-ylmethyl]-2-methylpropyl]-4-methylbenzamide dihydrochloride salt

[0182]



Step 1

[0183] To a suspension of 4-piperidone hydrochloride (3.52 g, 22.9 mmol) in ethanol (40 ml) was added triethylamine (10.8 ml, 91.6 mmol), followed by the addition of 3,4-dichlorobenzyl bromide (5.0 g, 20.8 mmol) and the reaction mixture was stirred overnight at room temperature. The reaction mixture was concentrated *in vacuo* and the residue was partitioned between water and ethyl acetate. The ethyl acetate layer was separated, washed with brine, dried over magnesium sulfate and concentrated *in vacuo*. Column chromatography with 25% ethyl acetate and 25% dichloromethane in hexanes gave 1-(3,4-dichlorobenzyl)-4-piperidone (4.4 g) as a colorless oil.

Step 2

[0184] To a suspension of sodium hydride (0.33 g, 13.2 mmol) in dry toluene (15 ml) was added triethyl phosphonoacetate (2.6 ml, 13.2 mmol) at a rate such that the internal reaction temperature remained below 25 °C. After stirring at room temperature for 0.5 h, the reaction mixture was recooled to below 25 °C and a solution of 1-(3,4-dichlorobenzyl)-4-piperidone (3.10 g, 12 mmol) in toluene (6 ml) was added dropwise. After the addition was complete, a gummy precipitate formed. The mixture was heated at 60-65 °C for 10 min. and then cooled to room temperature. The mother liquor was decanted and the residue was washed with more toluene. The combined toluene layers were filtered through a celite pad and concentrated *in vacuo* to give ethyl [1-(3,4-dichlorobenzyl)piperidin-4-ylidene]acetate (3.9 g) as an oil which was used in the next step without further purification.

Step 3

[0185] Ethyl [1-(3,4-dichlorobenzyl)piperidin-4-ylidene]acetate (3.9 g, 11.9 mmol) dissolved in tetrahydrofuran (25 ml) and ethanol (25 ml) was subjected to hydrogenation over platinum oxide (0.1 g) at ambient temperature and atmospheric pressure. The reaction mixture was filtered through a celite pad and concentrated *in vacuo* to give ethyl [1-(3,4-dichlorobenzyl)piperidin-4-yl]acetate (3.91 g) as a semi-solid.

Step 4

[0186] DIBAL-H (6.9 ml, 10.35 mmol, 1.5 M in toluene) was added to a solution of ethyl [1-(3,4-dichlorobenzyl)piperidin-4-yl]acetate (1.9 g, 5.8 mmol) in dry toluene (60 ml) at -60 °C under nitrogen atmosphere. The reaction mixture was stirred at dry ice bath temperature for 0.5 h, quenched with saturated sodium bisulfite solution and then allowed to warm to room temperature. The aqueous layer was separated and basified with 2N sodium hydroxide to pH between 8-9 and the product was extracted with diethyl ether. The combined ether layers were washed with brine and dried over sodium sulfate. After concentration, the residue was dissolved in dry tetrahydrofuran (30 ml) and cooled to 0 °C. Isopropyl magnesium chloride (2.0 M, 5.8 ml) was added dropwise and the resulting reaction mixture was allowed to warm to room temperature. After 0.5h, the reaction mixture was quenched with water and extracted with diethyl ether. The ether layer was washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified on silica gel column with 25% acetone, 25% dichloromethane in hexanes to give 1(*RS*)-[1-(3,4-dichlorobenzyl)piperidin-4-ylmethyl]-2-methylpropanol (0.75 g) as an oil.

Step 5

[0187] Pyridinium dichromate (4.0 g 10.63 mmol) was added to a solution of 1(*RS*)-[1-(3,4-dichlorobenzyl)piperidin-4-ylmethyl]-2-methylpropanol (0.5 g, 1.5 mmol) in dimethylformamide (25 ml) at 0 °C. The resulting solution was stirred at 0 °C until the starting material was consumed and then quenched with water, basified with 2N sodium hydroxide and extracted with ethyl acetate. The organic layer was washed with water, brine, and dried over sodium sulfate. The residue was purified on silica gel with 25% acetone, 25% dichloromethane in hexanes to give 1-[1-(3,4-dichlorobenzyl)piperidin-4-yl]-3-methyl-2-butanone (0.32 g) as an oil.

Step 6

[0188] To a solution of 1-[1-(3,4-dichlorobenzyl)piperidin-4-yl]-3-methyl-2-butanone (0.16 g, 0.49 mmol) in methanol (1.5 ml) was added ammonium acetate (0.38 g, 4.93 mmol), followed by sodium cyanoborohydride (31 mg, 0.5 mmol) and the resulting solution was stirred at room temperature. After 48 h, hydrochloric acid (conc.) was added until pH was 2 and the volatiles were removed under vacuum. The residue was dissolved in water and washed with ethyl acetate. The aqueous layer was then basified to pH>10 and extracted with ethyl acetate. The ethyl acetate layer was washed with brine, dried over sodium sulfate and concentrated to give 1(*RS*)-[1-(3,4-dichlorobenzyl)piperidin-4-ylmethyl]-2-methylpropylamine (0.16 g) as an oil which was used in the next step without further purification.


Step 7

[0189] *p*-Toluoyl chloride (0.077 ml, 0.58 mmol) was added to a solution of 1(*RS*)-[1-(3,4-dichlorobenzyl)piperidin-4-ylmethyl]-2-methylpropylamine (0.16 g, 0.49 mmol) in dichloromethane (2 ml) and triethylamine (0.12 ml, 1.63 mmol). The reaction mixture was stirred at room temperature for 1.5 h and was quenched with water. The product was then extracted with dichloromethane and the organic layer was washed with brine and dried over sodium sulfate. The crude product was purified on preparative TLC with 25% acetone, 25% dichloromethane in hexanes to give 0.21 g of N-[1-(1(*RS*)-[1-(3,4-dichlorobenzyl)piperidin-4-ylmethyl]-2-methylpropyl)-4-methylbenzamide as an oil, which was then converted to its dihydrochloride salt (0.15 g); m.p. 233.9-235.5 °C.

Example 11

N-{1-(*R*)-[4-(Dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl}-5-methylthiophene-2-carboxamide hydrochloride salt

[0190]




[0191] To a solution of 1-(*R*)-[4-(3,4-dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropylamine (0.52 g, 1.58 mmol) [prepared as described in Example 3, Step 5], in methylene chloride (15 ml) was added 5-methyl-2-thiophenecarboxylic acid (0.29 g, 2.05 mmol) followed by 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.39 g, 2.05 mmol), and the solution was stirred under argon overnight at room temperature. After concentrating the reaction mixture, water (10 ml) and ethyl acetate (25 ml) were added and the reaction mixture was stirred while adjusting to pH 8 with dilute aqueous potassium carbonate solution. The organic layer was separated and the aqueous layer was extracted with ethyl acetate (25 ml). The combined organic layers were dried over magnesium sulfate, filtered and concentrated *in vacuo*. The crude material (0.86g) was flash chromatographed on silica, eluting with 2% methanol/methylene chloride (containing 1% ammonium hydroxide) to give 0.64g product as a gum. Anhydrous HCl/ether (5 ml) was added to an ether solution of the product, giving a gummy precipitate which was further dissolved in methanol (10 ml) and stripped to give N-{1-(*R*)-[4-(dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl}-5-methylthiophene-2-carboxamide hydrochloride salt (0.72 g) as a light yellow foam.

Example 12

N-{1-(*R*)-[4-(Dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl}-4-(2-aminoethyl)benzamide dihydrochloride salt

[0192]



Step 1

[0193] A solution of 4-bromophenethylamine (10 g, 50 mmol), di-*tert*-butyldicarbonate (21.82 g, 100 mmol), triethylamine (13.9 ml, 100 mmol), and anhydrous methanol (350 ml) was refluxed for 2 h. The reaction mixture was concentrated *in vacuo* and the residue was stirred with water, filtered, washed with water and air dried to give N-(*tert*-butoxycarbonyl)-4-bromophenethylamine (15 g).

Step 2

[0194] To a 0 °C solution of N-(*tert*-butoxycarbonyl)-4-bromophenethylamine (12.82 g, 42.6 mmol) in anhydrous ether (200 ml) under argon was added dropwise *n*-butyllithium (34 ml, 85.2 mmol, 2.5 M in hexane). The slightly yellow solu-

tion turned milky white and was stirred 15 min., at 0° followed by one hour at room temperature. The resulting mixture was cooled to -78 °C in a dry ice-acetone bath. Dry carbon dioxide was bubbled through the stirring reaction mixture for 30 min., and then the reaction mixture was allowed to slowly warm to room temperature over 30 min. Water (100 ml) was added, the pH adjusted to 8 with dilute aqueous potassium carbonate solution, and the aqueous layer was separated and extracted with ethyl acetate. The aqueous layer was then cooled to 0 °C and acidified to pH 3 with 3 N aqueous hydrochloric acid. The resulting white precipitate was filtered, washed with water and air dried to provide 4-[2-(*N*-*tert*-butoxycarbonylamino)ethyl]benzoic acid (7.8 g).

Step 3

[0195] To a room temperature solution of 1-(*R*)-[4-(3,4-dichlorobenzyl)piperidin-1-yl-methyl]-2-methylpropylamine (1g, 3.04 mmol) (prepared as described in Example 3, Step 5) in methylene chloride (30 ml) was added 4-[2-(*N*-*tert*-butoxycarbonylamino)-ethyl]benzoic acid (1 g, 3.8 mmol) followed by 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.73 g, 3.8 mmol) and the solution was stirred under argon overnight at room temperature. After concentrating the reaction mixture, water (25 ml) and ethyl acetate (60 ml) were added and the reaction mixture was stirred while adjusting the pH to 8 with dilute aqueous potassium carbonate solution. The organic layer was separated and the aqueous layer was extracted with ethyl acetate (50 ml). The combined organic layers were dried over magnesium sulfate and concentrated. The resulting crude product was flash chromatographed on silica, eluting with 2%-4% methanol/methylene chloride containing 1% ammonium hydroxide to give *N*-{1-[1-(*R*)-[1-(3,4-dichlorobenzyl)-piperidin-4-ylmethyl]-2-methylpropyl]-4-[2-(*N*-*tert*-butoxycarbonylamino)ethyl]-benzamide as a white foam.


Step 4

[0196] To a room temperature solution of give *N*-{1-[1-(*R*)-[1-(3,4-dichlorobenzyl)-piperidin-4-ylmethyl]-2-methylpropyl]-4-[2-(*N*-*tert*-butoxycarbonylamino)ethyl]-benzamide (1.34 g, 2.32 mmol) in methylene chloride (30 ml) was added trifluoroacetic acid (30 ml). After 3 h, the reaction mixture was concentrated *in vacuo* and the residue was stirred with ethyl acetate (100 ml) and water (50 ml) while adjusting the pH to 8 with 15% aqueous sodium hydroxide solution. The organic layer was separated and the aqueous layer was extracted twice with ethyl acetate. The combined organic portions were dried with magnesium sulfate and concentrated. The crude was flash chromatographed on silica, eluting with 3% methanol/methylene chloride (containing 1% ammonium hydroxide). The free base product was dissolved in anhydrous ether and 1N anhydrous HCl / ether was added to give *N*-{1-(*R*)-[4-(dichlorobenzyl)piperidin-1-yl-methyl]-2-methylpropyl}-4-(2-aminoethyl)benzamide as the hydrochloride salt, mp 190.5-261.6 °C.

Example 13

1-{1-(*R*)-[4-(3,4-Dichlorobenzyl)piperidin-1-yl]-2-methylpropyl}-3-(3-methoxyphenyl)urea

[0197]



[0198] To a room temperature solution of 1-(*R*)-[4-(3,4-dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropylamine (0.349g, 1.06 mmol) (prepared as in Example 3, steps 4 and 5, but using *D*-Boc-Valine in place of *L*-Boc-Valine) in methylene chloride (50 ml) was added 3-methoxyphenylisocyanate (0.17ml, 1.27 mmol) and the reaction mixture was stirred at room temperature under argon for 2 h. The reaction mixture was concentrated *in vacuo* and the crude product was flash chromatographed on silica, eluting with 1-3% methanol/methylene chloride (containing 1% ammonium hydroxide). The product was dissolved in anhydrous ether and 1N anhydrous HCl / ether was added to precipitate 1-{1-(*R*)-[4-(3,4-dichlorobenzyl)piperidin-1-yl]-2-methylpropyl}-3-(3-methoxyphenyl)urea as the hydrochloride salt (0.29g), mp 107-112 °C.

Example 14

Formulation Examples

5 **[0199]** The following are representative pharmaceutical formulations containing a compound of Formula (I).

Tablet formulation

10 **[0200]** The following ingredients are mixed intimately and pressed into single scored tablets.

Ingredient	Quantity per tablet, mg
compound of this invention	400
cornstarch	50
croscarmellose sodium	25
lactose	120
magnesium stearate	5

Capsule formulation

25 **[0201]** The following ingredients are mixed intimately and loaded into a hard-shell gelatin capsule.

Ingredient	Quantity per capsule, mg
compound of this invention	200
lactose, spray-dried	148
magnesium stearate	2

Suspension formulation

The following ingredients are mixed to form a suspension for oral administration.

Ingredient	Amount
compound of this invention	1.0 g
fumaric acid	0.5 g
sodium chloride	2.0 g
methyl paraben	0.15 g
propyl paraben	0.05 g
granulated sugar	25.5 g
sorbitol (70% solution)	12.85 g
Veegum K (Vanderbilt Co.)	1.0 g
flavoring	0.035 ml
colorings	0.5 mg

(continued)

Ingredient	Amount
distilled water	q.s. to 100 ml

Injectable formulation

The following ingredients are mixed to form an injectable formulation.

Ingredient	Amount
compound of this invention	0.2 g
sodium acetate buffer solution, 0.4 M	2.0 ml
HCl (1N) or NaOH (1N)	q.s. to suitable pH
water (distilled, sterile)	q.s. to 20 ml

Example 15

CCR-3 Receptor Binding Assay- *in vitro*

[0202] The CCR-3 antagonistic activity of the compounds of the invention was determined by their ability to inhibit the binding of ^{125}I eotaxin to CCR-3 L1.2 transfectant cells ((see Ponath, P.D. et al., *J. Exp. Med.*, Vol. 183, 2437-2448, (1996)).

[0203] The assay was performed in Costar 96-well polypropylene round bottom plates. Test compounds were dissolved in DMSO and then diluted with binding buffer (50 mM HEPES, 1 mM CaCl_2 , 5 mM MgCl_2 , 0.5% bovine serum albumin (BSA), 0.02% sodium azide, pH 7.24) such that the final DMSO concentration was 2%. 25 μl of the test solution or only buffer with DMSO (control samples) was added to each well, followed by the addition of 25 μl of ^{125}I -eotaxin (100 pmol) (NEX314, New England Nuclear, Boston, MA) and 1.5×10^5 of the CCR-3 L1.2 transfected cells in 25 μl binding buffer. The final reaction volume was 75 μl .

[0204] After incubating the reaction mixture for 1h at room temperature, the reaction was terminated by filtering the reaction mixture through polyethylenimine treated Packard Unifilter GF/C filter plate (Packard, Chicago, IL.). The filters were washed four times with ice cold wash buffer containing 10 mM HEPES and 0.5M sodium chloride (pH 7.2) and dried at 65 °C for approximately 10 min. 25 μl /well of Microscint-20™ scintillation fluid (Packard) was added and the radioactivity retained on the filters was determined by using the Packard TopCount™.

[0205] Compounds of this invention were active in this assay.

[0206] The IC_{50} value (concentration of test compound required to reduce ^{125}I -eotaxin binding to the CCR-3 L 1.2 transfected cells by 50%) for some of the compounds of the invention was:

Cpd #	IC_{50} , μM	Cpd #	IC_{50} , μM
7	0.415	88	1.47
8	0.24	89	1.68
19	0.945	91	1.89
20	0.952	93	3.37
23	1.33	99	2.36
24	1.37	100	2.47
25	1.46	107	1.14
59	0.95	109	2.84

(continued)

Cpd #	IC ₅₀ , μ M	Cpd #	IC ₅₀ , μ M
61	1.76	110	3.05
78	5.03	117	1.33
84	0.69	118	1.5
86	1.16	121	3.52

Example 16

Inhibition of Eotaxin mediated chemotaxis of CCR-3 L1.2 transfectant cells--*In vitro* Assay

[0207] The CCR-3 antagonistic activity of the compounds of this invention was determined by measuring the inhibition of eotaxin mediated chemotaxis of the CCR-3 L1.2 transfectant cells, using a slight modification of the method described in Ponath, P. D. et al., *J. Clin. Invest.* 97: 604-612 (1996). The assay was performed in a 24-well chemotaxis plate (Costar Corp., Cambridge MA). CCR-3 L1.2 transfectant cells were grown in culture medium containing RPMI 1640, 10% Hyclone™ fetal calf serum, 55 mM 2-mercaptoethanol and Geneticin 418 (0.8 mg/ml). 18-24 hours before the assay, the transfected cells were treated with n-butyric acid at a final concentration of 5 mM/1 x 10⁶ cells/ml, isolated and resuspended at 1 x 10⁷ cells/ml in assay medium containing equal parts of RPMI 1640 and Medium 199 (M 199) with 0.5% bovine serum albumin.

[0208] Human eotaxin suspended in phosphate buffered saline at 1 mg/ml was added to bottom chamber in a final concentration of 100 nm. Transwell culture inserts (Costar Corp., Cambridge MA) having 3 micron pore size were inserted into each well and L1.2 cells (1 x 10⁶) were added to the top chamber in a final volume of 100 μ l. Test compounds in DMSO were added both to the top and bottom chambers such that the final DMSO volume was 0.5%. The assay was performed against two sets of controls. The positive control contained cells with no test compound in the top chamber and only eotaxin in the lower chamber. The negative control contained cells with no test compound in the top chamber and neither eotaxin nor test compound in lower chamber. The plate was incubated at 37 °C. After 4 h, the inserts were removed from the chambers and the cells that had migrated to the bottom chamber were counted by pipetting out 500 μ l of the cell suspension from the lower chamber to 1.2 ml Cluster tubes (Costar) and counting them on a FACS for 30 sec.

[0209] Compounds of this invention were active in this assay.

[0210] The IC₅₀ value (concentration of test compound required to reduce eotaxin mediated chemotaxis of CCR-3 L1.2 transfected cells by 50%) for some of the compounds of the invention was:

CPD#	IC ₅₀ μ M	CPD#	IC ₅₀ μ M
57	0.24	84	0.46
59	0.21	93	0.06

Example 17

Inhibition of Eotaxin mediated chemotaxis of human eosinophils--*In vitro* Assay

[0211] The ability of compounds of the invention to inhibit eotaxin mediated chemotaxis of human eosinophils was assessed using a slight modification of procedure described in Carr, M.W. et al., *Proc. Natl. Acad. Sci. USA*, 91: 3652-3656 (1994). Experiments were performed using 24 well chemotaxis plates (Costar Corp., Cambridge MA). Eosinophils were isolated from blood using the procedure described in PCT Application, Publication No. WO 96/22371. The endothelial cells used were the endothelial cell line ECV 304 obtained from European Collection of Animal Cell Cultures (Porton Down, Salisbury, U.K.). Endothelial cells were cultured on 6.5 mm diameter Biocoat® Transwell tissue culture inserts (Costar Corp., Cambridge MA) with a 3.0 μ M pore size. Culture media for ECV 304 cells consisted of M199, 10% Fetal Calf Serum, L-glutamine and antibiotics. Assay media consisted of equal parts RPMI 1640 and M199, with 0.5% BSA. 24 h before the assay 2 x 10⁵ ECV 304 cells were plated on each insert of the 24-well chemotaxis plate and incubated at 37 °C. 20 nM of eotaxin diluted in assay medium was added to the bottom chamber. The final volume in

bottom chamber was 600 μ l. The endothelial coated tissue culture inserts were inserted into each well. 10^6 eosinophil cells suspended in 100 μ l assay buffer were added to the top chamber. Test compounds dissolved in DMSO were added to both top and bottom chambers such that the final DMSO volume in each well was 0.5%. The assay was performed against two sets of controls. The positive control contained cells in the top chamber and eotaxin in the lower chamber. The negative control contained cells in the top chamber and only assay buffer in the lower chamber. The plates were incubated at 37 °C in 5% CO₂/95% air for 1-1.5 h.

[0212] The cells that had migrated to the bottom chamber were counted using flow cytometry. 500 μ l of the cell suspension from the lower chamber was placed in a tube, and relative cell counts were obtained by acquiring events for a set time period of 30 seconds.

[0213] Compounds of this invention were active in this assay.

Example 18

Inhibition of Eosinophil influx into the lungs of Ovalbumin sensitized balb/c mice by CCR-3 Antagonist -- *In vivo* Assay

[0214] The ability of the compounds of the invention to inhibit leukocyte infiltration into the lungs was determined by measuring the inhibition of eosinophil accumulation into the bronchioalveolar lavage (BAL) fluid of Ovalbumin (OA)-sensitized balb/c mice after antigen challenge by aerosol. Briefly, male balb/c mice weighing 20-25g were sensitized with OA (10 μ g in 0.2 ml aluminum hydroxide solution) intraperitoneally on days 1 and 14. After a week, the mice were divided into ten groups. Test compound or only vehicle (control group) or anti-eotaxin antibody (positive control group) was administered. After 1 h, the mice were placed in a Plexiglass box and exposed to OA aerosol generated by a PARISTAR™ nebulizer (PARI, Richmond, VA) for 20 min. Mice which had not been sensitized or challenged were included as negative control. After 24 or 72 h, the mice were anesthetized (urethane, approx. 1g/kg, i.p.), a tracheal cannula (PE 60 tubing) was inserted and the lungs were lavaged four times with 0.3 ml PBS. The BAL fluid was transferred into plastic tubes and kept on ice. Total leukocytes in a 20 μ l aliquot of the BAL fluid was determined by Coulter Counter™ (Coulter, Miami, FL). Differential leukocyte counts were made on Cytospin™ preparations which had been stained with a modified Wright's stain (Diff-Quick™) by light microscopy using standard morphological criteria.


[0215] Compounds of this invention were active in this assay.

[0216] The foregoing invention has been described in some detail by way of illustration and example, for purposes of clarity and understanding. It will be obvious to one of skill in the art that changes and modifications may be practiced within the scope of the appended claims. Therefore, it is to be understood that the above description is intended to be illustrative and not restrictive. The scope of the invention should, therefore, be determined not with reference to the above description, but should instead be determined with reference to the following appended claims, along with the full scope of equivalents to which such claims are entitled.

[0217] All patents, patent applications and publications cited in this application are hereby incorporated by reference in their entirety for all purposes to the same extent as if each individual patent, patent application or publication were so individually denoted.

Claims

1. A compound selected from compounds of Formula (I):



wherein:

T and U are both nitrogen; or one of T and U is nitrogen and the other is carbon;

R¹ and R² are, independently of each other, hydrogen or alkyl;

n is an integer from 0 to 2, provided that when n is 0, either T or U is carbon;

m is an integer from 0 to 3;

Ar and Ar¹ are, independently of each other, aryl or heteroaryl;

F is alkylene, alkenylene or a bond, provided that when T and U are nitrogen and F is alkylene, then R⁴ is not aryl.

Each R is independently hydrogen or alkyl, or R together with either R³ or R⁴ and the atoms to which they are attached form a carbocycle or a heterocycle;

R³ and R⁴ are, independently of each other, selected from:

(i) hydrogen, alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkylalkyl, heteroaryl, heteroaralkyl, heterocyclyl, heterocyclalkyl, heteroalkyl, cyano or

-(alkylene)-C(O)-Z where Z is alkyl, haloalkyl, alkoxy, haloalkyloxy, hydroxy, amino, mono- or disubstituted amino, aryl, aralkyl, aryloxy, aralkyloxy, heteroaryl, heteroaryloxy or heteroaralkyloxy, provided that both R³ and R⁴ are not hydrogen; or

(ii) R³ and R⁴ together with the carbon atom to which they are attached form a carbocycle or a heterocycle;

E is -C(O)N(R⁵)-, -SO₂N(R⁵)-, -N(R⁶)C(O)N(R⁵)-, -N(R⁶)SO₂N(R⁵)-, -N(R⁶)C(S)N(R⁵)-, -N(R⁶)C(O)-, -N(R⁶)C(O)O-, -OC(O)N(R⁶)- or -N(R⁶)SO₂- wherein:

R⁵ is:

(i) hydrogen, alkyl, acyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, heterocyclalkyl, heteroalkyl, or

-(alkylene)-C(O)-Z where Z is alkyl, haloalkyl, alkoxy, haloalkyloxy, hydroxy, amino, mono- or disubstituted amino, aryl, aralkyl, aryloxy, aralkyloxy, heteroaryl, heteroaryloxy or heteroaralkyloxy; or

(ii) R⁵ together with either R³ or R⁴ and the atoms to which they are attached forms a heterocycloamino group; and

R⁶ is hydrogen, alkyl, acyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, heterocyclalkyl, heteroalkyl, or -(alkylene)-C(O)-Z where Z is alkyl, haloalkyl, alkoxy, haloalkyloxy, hydroxy, amino, mono- or disubstituted amino, aryl, aralkyl, aryloxy, aralkyloxy, heteroaryl, heteroaryloxy or heteroaralkyloxy,

provided that when T is nitrogen and E is -C(O)N(R⁵)-, -SO₂N(R⁵)-, -N(R⁶)C(O)N(R⁵)-, -N(R⁶)SO₂N(R⁵)- or -N(R⁶)C(S)N(R⁵)-, then m > 0;

Q is -R⁷-W-R⁸- wherein:

R⁷ is an alkylene chain of between 1-6 carbon atoms inclusive;

R⁸ is a bond or an alkylene chain of between 0-4 carbon atoms inclusive;

W is a bond or a group selected from -C(O)-, -NR⁹-, -O-, -S(O)_{0.2}-, -C(O)N(R⁹)-, -N(R⁹)C(O)-, -N(R⁹)SO₂-, -SO₂N(R⁹)-, -N(R⁹)C(O)N(R⁹)-, -N(R⁹)SO₂N(R⁹)- or -N(R⁹)C(S)N(R⁹)- wherein: R⁹ is hydrogen, alkyl, acyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, aralkenyl, heteroaryl, heteroaralkyl, heterocycloalkyl, heteroalkyl, or -(alkylene)-C(O)-Z where Z is alkyl, haloalkyl, alkoxy, haloalkyloxy, hydroxy, amino, mono- or disubstituted amino, aryl, aralkyl, aryloxy, aralkyloxy, heteroaryl, heteroaryloxy or heteroaralkyloxy,

provided that when T is nitrogen and U is carbon then W is not -C(O)N(R⁹)-; and prodrugs, individual isomers, mixtures of isomers and pharmaceutically acceptable salts thereof.

2. The compound of Claim 1, wherein:

n and m are 1;

F is a bond;

Q is an alkylene chain of between 1 to 6 carbon atoms inclusive; and

E is -C(O)N(R⁵)-, -SO₂N(R⁵)-, -N(R⁶)C(O)N(R⁵)- or N(R⁶)C(O)-.

3. The compound of Claim 2, wherein:

R, R¹, R² and R³ are hydrogen; and

E is $-C(O)N(R^5)-$.

4. The compound of Claim 3, wherein T and U are both nitrogen.

5 5. The compound of Claim 4, wherein R^4 is alkyl or heteroalkyl and R^5 is hydrogen.

6. The compound of Claim 5, wherein:

Ar and Ar^1 are aryl.

10

7. The compound of Claim 6, wherein R^4 is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl.

8. The compound of Claim 7, wherein:

15

Ar is a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, $-SO_2R$ (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, $-CONR'R''$ (where R' and R'' are hydrogen or alkyl) or $-COOH$; and Ar^1 is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino.

20

9. The compound of Claim 8, wherein:

Ar is phenyl, 4-chlorophenyl, 3,4-difluorophenyl, 4-methylphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 4-hydroxyphenyl, 3,4-methylenedioxyphenyl, 4-methylsulfonylphenyl, 4-[(2-acetylamino)ethyl]phenyl, 4-{2-[(*R*)-amino-3-methylbutylamino]ethyl}phenyl, 4-(2-aminoethyl)phenyl, 4-(aminomethyl)-phenyl, 4-(hydroxymethyl)phenyl, 3-aminocarbonylphenyl, 3-carboxyphenyl, 2,5-dimethoxyphenyl, 3,5-dimethoxyphenyl, 3,4-dimethoxyphenyl or 4-aminophenyl; and Ar^1 is 4-nitrophenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 3,4-difluorophenyl, 2,3-dichlorophenyl, 3-methyl-4-nitrophenyl, 3-chloro-4-fluoro-phenyl or 3,4-dichlorophenyl.

30

10. The compound of Claim 9, wherein:

R^4 is 1-methylethyl;
Ar is 4-methylphenyl;
 Ar^1 is 3,4-dichlorophenyl; and
Q is methylene;

35

namely, N-{1-(*S*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-4-methylbenzamide dihydrochloride salt.

40

11. The compound of Claim 9, wherein:

R^4 is 1,1-dimethylethyl;
Ar is 4-methylphenyl;
 Ar^1 is 3,4-dichlorophenyl; and
Q is methylene;

45

namely, N-{1-(*S*)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2,2-dimethylpropyl}-4-methylbenzamide dihydrochloride salt.

50

12. The compound of Claim 5, wherein:

Ar is an aryl ring; and
 Ar^1 is a heteroaryl ring.

55

13. The compound of Claim 12, wherein R^4 is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl.

14. The compound of Claim 13, wherein:

Ar is a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH; and Ar¹ is 1-acetylindol-3-yl, 3-methylbenzothiophen-2-yl or 5-nitrothiophen-3-yl.

15. The compound of Claim 5, wherein:

Ar is a heteroaryl ring; and
Ar¹ is an aryl ring.

16. The compound of Claim 15, wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl.

17. The compound of Claim 16, wherein:

Ar is pyridin-2-yl, pyridin-3-yl, quinolin-3-yl or 5-methylthiophen-2-yl;
and
Ar¹ is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino.

18. The compound of Claim 17, wherein:

Ar¹ is 4-nitrophenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 3,4-difluorophenyl, 2,3-dichlorophenyl, 3-methyl-4-nitrophenyl, 3-chloro-4-fluorophenyl or 3,4-dichlorophenyl.

19. The compound of Claim 3, wherein T is nitrogen and U is carbon.

20. The compound of Claim 19, wherein R⁴ is alkyl or heteroalkyl and R⁵ is hydrogen.

21. The compound of Claim 20, wherein:

Ar and Ar¹ are aryl.

22. The compound of Claim 21, wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl.

23. The compound of Claim 22, wherein:

Ar is a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH; and
Ar¹ is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino.

24. The compound of Claim 23, wherein:

Ar is phenyl, 4-chlorophenyl, 3,4-difluorophenyl, 4-methylphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 4-hydroxyphenyl, 3,4-methylenedioxyphenyl, 4-methylsulfonylphenyl, 4-[(2-acetylamino)ethyl]phenyl, 4-{2-[(R)-amino-3-methylbutyrylamino]ethyl}phenyl, 4-(2-aminoethyl)phenyl, 4-(aminomethyl)-phenyl, 4-(hydroxymethyl)phenyl, 3-aminocarbonylphenyl, 3-carboxyphenyl, 2,5-dimethoxyphenyl, 3,5-dimethoxyphenyl, 3,4-dimethoxyphenyl or 4-aminophenyl; and
Ar¹ is 4-nitrophenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 3,4-difluorophenyl, 2,3-dichlorophenyl, 3-methyl-4-nitrophenyl, 3-chloro-4-fluoro-phenyl or 3,4-dichlorophenyl.

25. The compound of Claim 24, wherein:

R⁴ is 1-methylethyl;
 Ar is 4-methylphenyl;
 Ar¹ is 3,4-dichlorophenyl; and
 Q is methylene;

and is named as, N-{1-(*S*)-[4-(3,4-dichlorobenzyl)piperidin-1-ylmethyl]-2-methyl-propyl}-4-methylbenzamide dihydrochloride salt.

26. The compound of Claim 24, wherein:

R⁴ is 1-methylethyl;
 Ar is 4-(2-aminoethyl)phenyl;
 Ar¹ is 3,4-dichlorophenyl; and
 Q is methylene;

and is named as, N-{1-(*R*)-[4-(3,4-dichlorobenzyl)piperidin-1-ylmethyl]-2-methyl-propyl}-4-(2-aminoethyl)benzamide dihydrochloride salt.

27. The compound of Claim 20, wherein:

Ar is a heteroaryl ring; and
 Ar¹ is an aryl.

28. The compound of Claim 27, wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl.

29. The compound of Claim 28, wherein:

Ar is pyridin-2-yl, pyridin-3-yl, quinolin-3-yl or 5-methylthiophen-2-yl;
 and
 Ar¹ is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino.

30. The compound of Claim 29, wherein:

Ar¹ is 4-nitrophenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 3,4-difluorophenyl, 2,3-dichlorophenyl, 3-methyl-4-nitrophenyl, 3-chloro-4-fluoro-phenyl or 3,4-dichlorophenyl.

31. The compound of Claim 31 wherein:

R⁴ is 1-methylethyl;
 Ar is 5-methylthiophen-2-yl;
 Ar¹ is 3,4-dichlorophenyl; and
 Q is methylene;

and is named as, N-{1-(*R*)-[4-(3,4-dichlorobenzyl)piperidin-1-ylmethyl]-2-methyl-propyl}-5-methylthiophene-2-carboxamide hydrochloride salt.

32. The compound of Claim 3, wherein T is carbon and U is nitrogen.

33. The compound of Claim 32, wherein R⁴ is alkyl or heteroalkyl; and R⁵ is hydrogen.

34. The compound of Claim 33, wherein:

Ar is an aryl or heteroaryl ring; and
 Ar¹ is an aryl ring.

35. The compound of Claim 34, wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-

hydroxyethyl or 2-hydroxyethyl.

36. The compound of Claim 35, wherein:

Ar is a pyridin-2-yl, pyridin-3-yl, quinolin-3-yl or 5-methylthiophen-2-yl ring, or a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH; and
Ar¹ is a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino.

37. The compound of Claim 2, wherein:

R, R¹, R² and R³ are hydrogen; and
E is -N(R⁶)C(O)N(R⁵)-

38. The compound of Claim 37, wherein T and U are both nitrogen.

39. The compound of Claim 38, wherein R⁴ is alkyl or heteroalkyl; and R⁵ and R⁶ are hydrogen.

40. The compound of Claim 39, wherein:

Ar and Ar¹ are aryl.

41. The compound of Claim 40, wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl.

42. The compound of Claim 41, wherein:

Ar is a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH; and
Ar¹ is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino.

43. The compound of Claim 42, wherein:

Ar is phenyl, 4-chlorophenyl, 3,4-difluorophenyl, 4-methylphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 4-hydroxyphenyl, 3,4-methylenedioxyphenyl, 4-methylsulfonylphenyl, 4-[(2-acetylamino)ethyl]phenyl, 4-{2-[(R)-amino-3-methylbutyrylamino]ethyl}phenyl, 4-(2-aminoethyl)phenyl, 4-(aminomethyl)-phenyl, 4-(hydroxymethyl)phenyl, 3-aminocarbonylphenyl, 3-carboxyphenyl, 2,5-dimethoxyphenyl, 3,5-dimethoxyphenyl, 3,4-dimethoxyphenyl or 4-aminophenyl; and
Ar¹ is 4-nitrophenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 3,4-difluorophenyl, 2,3-dichlorophenyl, 3-methyl-4-nitrophenyl, 3-chloro-4-fluorophenyl or 3,4-dichlorophenyl.

44. The compound of Claim 43, wherein:

R⁴ is 1-methylethyl;
Ar is 3-methoxyphenyl;
Ar¹ is 3,4-dichlorophenyl; and
Q is methylene;

namely, 1-{1(R)-[4-(3,4-dichlorobenzyl)piperazin-1-ylmethyl]-2-methylpropyl}-3-(3-methoxyphenyl)urea.

45. The compound of Claim 37, wherein T is nitrogen and U is carbon.

46. The compound of Claim 45, wherein R⁴ is alkyl or heteroalkyl; and R⁵ and R⁶ are hydrogen.

47. The compound of Claim 46, wherein:

Ar and Ar¹ are aryl.

5 48. The compound of Claim 47, wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl.

49. The compound of Claim 48, wherein:

10 Ar is a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH; and Ar¹ is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino.

15 50. The compound of Claim 49, wherein:

20 Ar is phenyl, 4-chlorophenyl, 3,4-difluorophenyl, 4-methylphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 4-hydroxyphenyl, 3,4-methylenedioxyphenyl, 4-methylsulfonylphenyl, 4-[(2-acetylamino)ethyl]phenyl, 4-{2-[(R)-amino-3-methylbutyrylamino]ethyl}phenyl, 4-(2-aminoethyl)phenyl, 4-(aminomethyl)-phenyl, 4-(hydroxymethyl)phenyl, 3-aminocarbonylphenyl, 3-carboxyphenyl, 2,5-dimethoxyphenyl, 3,5-dimethoxyphenyl, 3,4-dimethoxyphenyl or 4-aminophenyl; and Ar¹ is 4-nitrophenyl, 4-trifluoromethylphenyl, 4-chlorophenyl, 3,4-difluorophenyl, 2,3-dichlorophenyl, 3-methyl-4-nitrophenyl, 3-chloro-4-fluorophenyl or 3,4-dichlorophenyl.

25 51. The compound of Claim 50, wherein:

30 R⁴ is 1-methylethyl;
Ar is 3-methoxyphenyl;
Ar¹ is 3,4-dichlorophenyl; and
Q is methylene;

namely, 1-{1-(R)-[4-(3,4-dichlorobenzyl)piperidin-1-ylmethyl]-2-methylpropyl}-3-(3-methoxyphenyl)urea.

35 52. The compound of Claim 37, wherein T is carbon and U is nitrogen.

53. The compound of Claim 52, wherein R⁴ is alkyl or heteroalkyl; and R⁵ and R⁶ are hydrogen.

40 54. The compound of Claim 53, wherein:

Ar and Ar¹ are aryl.

55 55. The compound of Claim 54, wherein R⁴ is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl.

56. The compound of Claim 55, wherein:

50 Ar is a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, -SO₂R (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, -CONR'R'' (where R' and R'' are hydrogen or alkyl) or -COOH; and Ar¹ is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino.

57. The compound of Claim 1, wherein:

55 T is carbon and U is nitrogen;
n is 1;
m is 0;

F is a bond;

Q is an alkylene chain; and

E is $-C(O)N(R^5)-$, $-SO_2N(R^5)-$, $-N(R^6)C(O)N(R^5)-$ or $N(R^6)C(O)-$.

5 58. The compound of Claim 57, wherein:

R, R^1 , R^2 and R^3 are hydrogen; and

E is $-C(O)N(R^5)-$.

10 59. The compound of Claim 58, wherein R^4 is alkyl or heteroalkyl; and R^5 is hydrogen.

60. The compound of Claim 59, wherein:

Ar is an aryl or heteroaryl ring; and

15 Ar^1 is an aryl ring.

61. The compound of Claim 60, wherein R^4 is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl.

20 62. The compound of Claim 61, wherein:

Ar is pyridin-2-yl, pyridin-3-yl, quinolin-3-yl or 5-methylthiophen-2-yl, or a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, $-SO_2R$ (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, $-CONR'R''$ (where R' and R'' are hydrogen or alkyl) or $-COOH$; and

25 Ar^1 is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino.

63. The compound of Claim 57, wherein:

30 R, R^1 , R^2 and R^3 are hydrogen; and
E is $-(NR^6)C(O)N(R^5)-$.

64. The compound of Claim 63, wherein R^4 is alkyl or heteroalkyl; and R^5 and R^6 are hydrogen.

35 65. The compound of Claim 64, wherein:

Ar is an aryl or heteroaryl ring; and

40 Ar^1 is an aryl ring.

66. The compound of Claim 65, wherein R^4 is 1-methylethyl, 1,1-dimethylethyl, 2-methylpropyl, 3-hydroxypropyl, 1-hydroxyethyl or 2-hydroxyethyl.


67. The compound of Claim 66, wherein:

45 Ar is pyridin-2-yl, pyridin-3-yl, quinolin-3-yl or 5-methylthiophen-2-yl, or a phenyl ring optionally substituted with one, two or three substituents selected from alkyl, heteroalkyl, alkoxy, acyl, $-SO_2R$ (where R is alkyl, amino or mono or disubstituted amino), methylenedioxy, hydroxy, halo, amino, mono- or disubstituted amino, $-CONR'R''$ (where R' and R'' are hydrogen or alkyl) or $-COOH$; and

50 Ar^1 is a phenyl ring optionally substituted with one, two or three substituent selected from alkyl, heteroalkyl, alkoxy, halo, trifluoromethyl, nitro or mono- or disubstituted amino.

68. A pharmaceutical composition comprising a therapeutically effective amount of a compound of any one of Claims 1 to 67 and a pharmaceutically acceptable excipient.


55 69. Compounds of the general formula:



15 in which n , m , R , $R^1 - R^4$, T , U , Q and Ar^1 are as in Claim 1 and X is a group $HN(R^5)$, $HN(R^6)$ - or carboxy, and R^5 and R^6 are as in Claim 1.

20 **70.** Process for the manufacture of the compounds in accordance with any one of Claims 1 - 67, which comprises

(a) transforming the group X in a compound of the general formula




35 in which n , m , R , $R^1 - R^4$, T , U , Q and Ar^1 are as in Claim 1 and X is a group $HN(R^5)$, $HN(R^6)$ - or carboxy, and R^5 and R^6 are as in Claim 1, into a group $Ar-(F)-(E)$ -, wherein Ar , F and E are as in Claim 1, or

(b) for the manufacture of a pharmaceutically acceptable salt of a compound of formula I carrying an acidic and/or basic substituent, converting such compound of formula I into such salt.

40 **71.** Compounds according to any one of Claims 1-67 whenever manufactured according to the process claimed in Claim 70 or by an obvious chemical equivalent thereof.

72. Compounds according to any one of Claims 1-67 for use as therapeutically active substances, against respiratory diseases such as asthma.

45 **73.** The use of compounds according to any one of Claims 1-67 in the control or prevention of diseases, particularly asthma, or for the manufacture of pharmaceutical preparations having such activity.



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 0 903 349 A3**

(12) **EUROPEAN PATENT APPLICATION**

(88) Date of publication A3:
24.05.2000 Bulletin 2000/21

(43) Date of publication A2:
24.03.1999 Bulletin 1999/12

(21) Application number: **98114971.9**

(22) Date of filing: **10.08.1998**

(51) Int. Cl.⁷: **C07D 295/12**, C07D 207/09,
C07D 207/20, C07D 209/42,
C07D 211/16, C07D 211/18,
C07D 213/81, C07D 215/48,
C07D 215/54, C07D 307/68,
C07D 307/84, C07D 317/68,
C07D 333/34, C07D 333/38,
C07D 333/68, C07D 401/12,
C07D 409/12, A61K 31/445,
A61K 31/495

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **18.08.1997 US 56001 P**

(71) Applicant:
**F. HOFFMANN-LA ROCHE AG
4070 Basel (CH)**


(72) Inventors:
• **Gong, Leyi**
San Mateo, California 94404 (US)

• **Kertesz, Denis John**
Mountain View, California 94043 (US)
• **Smith, David Bernard**
San Mateo, California 94403 (US)
• **Talamas, Francisco Xavier**
San Carlos, California 94070 (US)
• **Wilhelm, Robert Stephen**
Los Altos, California 94022 (US)

(74) Representative: **Braun, Axel et al**
F.Hoffmann-La Roche AG
Patent Department (PLP),
124 Grenzacherstrasse
4070 Basel (CH)

(54) **CCR-3 receptor antagonists**

(57) This invention relates to certain cyclic amine derivatives of Formula (I)



EP 0 903 349 A3

that are CCR-3 receptor antagonists, pharmaceutical compositions containing them, methods for their use and methods for preparing these compounds.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 11 4971

DOCUMENTS CONSIDERED TO BE RELEVANT					
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)		
A	EP 0 395 313 A (AMERICAN HOME PRODUCTS CORPORATION) 31 October 1990 (1990-10-31) * the whole document *	1,73	C07D295/12 C07D207/09 C07D207/20 C07D209/42		
A	EP 0 311 517 A (TAKEDA CHEMICAL INDUSTRIES, LTD) 12 April 1989 (1989-04-12) * the whole document *	1,73	C07D211/16 C07D211/18 C07D213/81 C07D215/48 C07D215/54		
A	EP 0 217 286 A (OKAMOTO SHOSUKE ET AL.) 8 April 1987 (1987-04-08) * the whole document *	1,73	C07D307/68 C07D333/38 C07D333/68 C07D307/84		
A	PATENT ABSTRACTS OF JAPAN vol. 13, no. 101 (C-574), 9 March 1989 (1989-03-09) & JP 63 275576 A (NIPPON CHEMIPHAR CO LTD), 14 November 1988 (1988-11-14) * abstract *	1,73	C07D317/68 C07D401/12 C07D409/12 C07D333/34 A61K31/445		
A	PATENT ABSTRACTS OF JAPAN vol. 13, no. 72 (C-570), 17 February 1989 (1989-02-17) & JP 63 264525 A (NIPPON CHEMIPHAR CO LTD), 1 November 1988 (1988-11-01) * abstract *	1,73	<table border="1"> <thead> <tr> <th>TECHNICAL FIELDS SEARCHED (Int.Cl.6)</th> </tr> </thead> <tbody> <tr> <td>C07D</td> </tr> </tbody> </table>	TECHNICAL FIELDS SEARCHED (Int.Cl.6)	C07D
TECHNICAL FIELDS SEARCHED (Int.Cl.6)					
C07D					
The present search report has been drawn up for all claims					
Place of search BERLIN		Date of completion of the search 5 April 2000	Examiner Kyriakakou, G		
<table border="0"> <tr> <td style="vertical-align: top;"> CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document </td> <td style="vertical-align: top;"> T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document </td> </tr> </table>				CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document	T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document	T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document				

EPO FORM 1503 03/82 (F04C01)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 11 4971

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
			A61K31/495
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 5 April 2000	Examiner Kyriakakou, G
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03/02 (P4/C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 11 4971

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

05-04-2000

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 395313 A	31-10-1990	AT 179973 T	15-05-1999
		AT 187718 T	15-01-2000
		AU 619677 B	30-01-1992
		AU 5377890 A	25-10-1990
		AU 619678 B	30-01-1992
		AU 5377990 A	25-10-1990
		CA 2015033 A	22-10-1990
		CA 2015034 A	22-10-1990
		DD 296921 A	19-12-1991
		DD 297968 A	30-01-1992
		DE 69033100 D	17-06-1999
		DE 69033100 T	05-01-2000
		DE 69033393 D	20-01-2000
		EP 0395312 A	31-10-1990
		EP 0955296 A	10-11-1999
		ES 2130116 T	01-07-1999
		FI 93832 B	28-02-1995
		FI 102175 B	30-10-1998
		GB 2230780 A,B	31-10-1990
		GB 2230781 A,B	31-10-1990
		GB 2255976 A,B	25-11-1992
		GR 3030667 T	29-10-1999
		HK 171395 A	17-11-1995
		HU 54667 A	28-03-1991
		HU 54666 A	28-03-1991
		HU 9500568 A	30-10-1995
		IE 65362 B	18-10-1995
		IE 64038 B	28-06-1995
		IL 94151 A	31-08-1995
		IL 94160 A	24-06-1994
		JP 3011059 A	18-01-1991
		JP 3020263 A	29-01-1991
		KR 142417 B	01-06-1998
		KR 173310 B	01-02-1999
		NZ 233398 A	23-12-1991
		NZ 233401 A	26-03-1991
		PH 26582 A	19-08-1992
		PT 93824 A,B	20-11-1990
		PT 93825 A,B	20-11-1990
		US 5340812 A	23-08-1994
		US 5420278 A	30-05-1995
		US 5541326 A	30-07-1996
		US 4921958 A	01-05-1990
		US 4988814 A	29-01-1991
		US 5364849 A	15-11-1994
		US 5382583 A	17-01-1995

EPC FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 11 4971

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

05-04-2000

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 395313 A		ZA 9003019 A	24-12-1991
		ZA 9003020 A	24-12-1991
EP 311517 A	12-04-1989	JP 1207284 A	21-08-1989
		US 4990614 A	05-02-1991
EP 217286 A	08-04-1987	AU 598750 B	05-07-1990
		AU 6305186 A	02-04-1987
		CA 1297633 A	17-03-1992
		US 4895842 A	23-01-1990
		JP 2023215 C	26-02-1996
		JP 7053705 B	07-06-1995
		JP 63022061 A	29-01-1988
JP 63275576 A	14-11-1988	JP 2536754 B	18-09-1996
JP 63264525 A	01-11-1988	JP 2515119 B	10-07-1996

EPO FORM P458

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82